

**Original Article****Dietary Patterns and Their Associations with General and Abdominal Obesity Among Adults in Gonabad in 2019: A Cross-Sectional Study**

Maryam Esmaeili¹, Mojtaba Kianmehr², Maryam Eskafi Noghani³, Amirhosein Basirimoghadam⁴, Mehrdad Kianmehr⁵, Mustafa Pouryousef⁶, Hamid Rasekhi^{7*}

1- MSc in Social Sciences, Gonabad University of Medical Sciences, Gonabad, Iran

2- Professor, Department of Medical Physics, Faculty of Medicine, Gonabad University of Medical Sciences, Gonabad, Iran

3- Assistant Professor, Department of Social Sciences, Faculty of Humanities, Islamic Azad University Gonabad Branch, Gonabad, Iran

4- Medical Student, Faculty of Medicine, Gonabad University of Medical Sciences, Gonabad, Iran

5- BSc Student in Nutrition Sciences, Department of Nutrition Sciences, Varastegan Institute for Medical Sciences, Mashhad, Iran

6- BSc Student in Educational Sciences, Department of Human Sciences, Shahid Rajaei Institute for Teacher Training, Torbat Heydariyeh, Iran

7- Department of Nutrition Research, Faculty of Nutrition and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Science, Tehran, Iran

Received: September 2020

Accepted: December 2020

ABSTRACT

Background and Objectives: Obesity is serious and most important underlying risk factor on incidence of non-communicable diseases (NCDs). In the last two decades, obesity has been on an increasing trend globally and across all age groups. Using dietary pattern analysis method could provide more information about nutritional etiology of chronic disease such as obesity. The aim of this study is to determine the association between major dietary patterns and general and central obesity among adults in Gonabad in 2019.

Materials and Methods: This cross-sectional study included 250 individuals aged 18-70 years living in Gonabad, Iran. Participants were selected by a multi-stage sampling method. The dietary intake of participants during the past year was evaluated using the semi-quantitative Food Frequency Questionnaire (FFQ), which included 168 foods items. Factor analysis was used for identifying major dietary patterns. The relationship between major dietary patterns and general and abdominal obesity was assessed by chi-square test. All data were analyzed by SPSS 14 at significance level of $P < 0.05$.

Results: Participants ages ranged between 18 and 70 years old, of whom 44.8% were male and 82.4% were married. Two healthy and unhealthy dietary patterns were identified. There was no significant relationship between unhealthy diet and height ($p=0.82$) and waist circumference ($p=0.32$) variables, but this dietary pattern was correlated with body weight ($p=0.003$), hip circumference ($p < 0.001$) and body mass index ($p < 0.001$). Also, there were significant relationship between healthy dietary pattern and height ($p < 0.001$), weight ($p < 0.001$), waist circumference ($p < 0.001$), hip circumference ($p < 0.001$) and body mass index ($p=0.001$). The unhealthy diet was significantly associated with general obesity ($p < 0.001$) and abdominal obesity ($p=0.01$). Both general and abdominal obesity measures were significantly associated with Healthy dietary pattern ($p = 0.003$ and $p < 0.001$ respectively).

Conclusions: Major dietary patterns were significantly associated with general and abdominal obesity among adults in Gonabad. Further studies with future prospective studies, are required to confirm these findings.

Keywords: Dietary patterns, Obesity, Adults, Gonabad, Iran

Introduction

Global evidence indicates that overweight and obesity are two of the main public health problems in all age groups in the population. The World Health Organization (WHO) report states that over the past four decades, the prevalence of overweight and obesity has increased dramatically in the general population. In 2016, 39 and 13% of adults in the world were overweight and obese, respectively(1).

The prevalence of obesity and central adiposity (enlarged waist circumference) has risen steeply in Iran, particularly among women, and is becoming a public health issue. The prevalence of obesity and overweight/obesity among Iranian adults is, 22.7% and 59.3% respectively. There is a significant difference between the prevalence of obesity among males 15.3% and females 29.8%(2).

Obesity is due to a number of factors including genetics, metabolic, socioeconomic status, environmental and behavioral factors (3). The underlying link between overweight, obesity and many chronic diseases such as cardiovascular disease, diabetes, rheumatoid arthritis, hypertension, physical inability and respiratory illness have strengthened the importance of these public health issues (1, 4, 5).

Diet is a component of lifestyle, which plays an important role in developing overweight and obesity (6, 7). Recently, the analysis of dietary pattern as an approach to study the links between food and disease was recommended. (8, 9).

In this approach, statistical methods are used to combine several foods or nutrients in order to derive single-exposure variables, or dietary patterns(10). It has been suggested that such dietary patterns could provide a better and more general overview of relation between diet and disease. (11) and perhaps more predictive of chronic disease risk than the intake of individual nutrients or foods(10).

Several studies have reported the association of the major dietary patterns with general and central obesity (12-18).

Also high prevalence of obesity in Iran and is associated with an elevated risk of several major non-communicable diseases(19), and knowing that food patterns are vary in different cultures, races, genders and geographical areas (20) and based on the fact that there has been no study of nutritional patterns in Gonabad so far, this study aims to determine the major dietary patterns and their relationship with obesity in adults people in Gonabad in 2019.

Given the high prevalence of non-communicable diseases related to nutrition, and also considering that dietary patterns in different cultures, races, sex and geographical regions are different, and since no studies have been conducted on dietary patterns in Gonabad City, this study aimed to determine dietary patterns and their relationship with obesity in adults.

Materials and Methods

This study was a cross-sectional analytical study. The study population consisted of male and female adults living in Gonabad in 2019. This research was reviewed and approved by the Ethics Committee in Gonabad University of Medical Sciences (ethical code: IR.GMU.REC.1397.060 and proposal cod: p/1/477). For the factor analysis, 24 food groups of dominant dietary patterns were studied based on Table1. For each variable least 10 people were needed(21, 22), so the sample size was set at 240 individuals. To account for the 10% dropout, the final sample size was set at 264. The Samples were selected using multistage random sampling, using a combination of simple,

systematic and random cluster sampling methods. The sampling method was tested on patient who referred to Gonabad health centers. Then based on the number of populations covered by that center, the required sample size was calculated and several clusters were selected. In each zone, based on the systematic sampling of each family, one eligible person in a family was randomly interviewed, and completed the questionnaires. Inclusion criteria were aged between 18 and 70 years, gonabad resident for at least 5 years., stable physiological and psychological status and farsi literate to answer questions. pregnant or breast-feeding women, people with disorders of the thyroid gland, and weight loss or obesity diet, were excluded(23). After obtaining written informed consent, the researcher collected demographic and nutritional information of the samples via interview and completing food frequency questionnaire. Those who did not respond to more than 50% of the food frequency questionnaire were excluded (24-26). After excluding those who did not answer more than 50% of the Food Frequency Questionnaire (FFQ), so 250 remained for the final analysis.

Dietary information was collected by the food frequency questionnaire containing 168 types of nutrients along with a standard amount of each nutrient for individuals. The participants were asked to report their consumption frequency of each food item in the past year by the type of food on a daily (e.g., bread), weekly (e.g., rice), or monthly (e.g., fish) basis. The values listed for each meal were calculated using the home-scale guide to grams, then the gram intake of each food per day was calculated for each individual(27). Food items are grouped into one factor based on their degree of correlation. Each score for a particular dietary pattern was calculated by multiplying the amount of each food consumed in that pattern. Then, the participants were classified according to the quota of dietary habits (27-31). In order to identify the dietary patterns and perform principal component analysis, due to the large number of items on the FFQ, the food items were first categorized to 24 food groups, based on the similarity of their nutrients and researchers' opinions, as well as on previous studies(32)

The weight was measured using the Seca 881 digital weighing scale to the nearest 0.1 kg without shoes while wearing minimal clothes. Height was measured to the nearest 1 mm without shoes with shoulders in a normal position. BMI (Body Mass Index) was calculated as weight in kilograms divided by height in meters squared. General obesity was defined as $BMI \geq 30 \text{ kg/m}^2$. Waist Circumference(WC) and hip Circumference(HC) were measured with an inelastic plastic tape measure according to WHO guidelines(33). Central obesity was defined as $WC \geq 88 \text{ cm}$ (34-38).

Table 1. Food groups used in the analysis of adult food patterns in Gonabad

	Food Groups	Food Items
1	Refined cereals	Lavash Bread, Baguette Bread, Starch, Wheat Flour, Rice, Macaroni, String, Vermicelli
2	Whole grains	Sangak bread, Taftoon bread, Barberry bread, barley bread, wheat
3	Potato	Potatoes (all preparation methods)
4	Tomato	Tomato, tomato juice, tomato paste, ketchup
5	Egg	Egg whites, egg yolks
6	Processed meats	Hot dogs, sausages, hamburger
7	Legumes	Lentils, cowpeas, beans, peas, cotyledons, soybeans
8	Nuts and seeds	Chickpea, Walnut, Hazelnut, Almond, Pistachio, Peanut, Indian Almond, Sesame, Seeds
9	Solid fats	Butter, Creamy, Creamy cheese, Skim, Chocolate Cheese, Pizza Cheese
10	Salt	Salt
11	Salty snacks	Salted crackers, pretzels, chips, snack, pickles
12	Jam and compote	All kinds of jams and compotes
13	Vegetables	Spinach, lettuce, Carrot, Yellow (winter) squash, cucumber, pepper, mushroom, onion, garlic, green beans, sweet corn, eggplant, mushrooms, celery, cabbage, Brussels sprouts, cauliflower, broccoli
14	Organ meats	Liver, heart, kidney, Chicken liver, gizzard, tripe, head, tongue, brain
15	Red meat	Beef, mutton, camel meat, hamburger
16	Mayonnaise	A variety of mayonnaise and salad dressings
17	Soft drinks	Industrial juices with added sugar, beverages, syrups, non-alcoholic beer
18	Coffee and tea	Tea and coffee
19	chicken and fish	Fish, Tuna, Chicken (all methods of preparation)
20	Fruits	Pears, apricots, cherries, apples, raisins or grapes, bananas, cantaloupe, watermelon, oranges, grapefruit, kiwi, strawberries, peaches, nectarine, tangerine, mulberry, plums, persimmons, pomegranates, lemons, pineapples, fresh figs, dates, dried figs, dried dates, other dried fruit and natural juices.
21	Dairy	Milk, Powdered Milk, Cocoa Milk, Yogurt, Cheese, Pasteurized Curd, Dough, Ice Creams
22	Vegetable oils	Sunflower oil, corn oil, olive oil and olive
23	Cookies	Cakes, muffins, Gaz, roulette, Danish cake
24	Sweet snacks	Sugar, Chocolate, Toffee, Sohan, Pastel, Jelly, Caramel Cream, Chocolate Biscuit, Creamy Biscuit, Wafer, Chewing gum

Demographic characteristics were age, sex, occupation, education marital status the and number of family members. The relationship between, major dietary patterns and demographic factors was assessed.

For statistical analysis, the obtained data were fed in SPSS v.14. The dominant dietary patterns (factors) were identified utilizing factor analysis. The participants were scored for each identified food pattern [25]. Factor scores were categorized as tertiles. Dietary pattern naming was conducted based on the interpretation of the food items per factor and according to previous studies [26]

Continuous and categorical variables were compared across groups using analysis of variance One-way analysis of variance (ANOVA) and Chi-square tests, respectively. The significance level of less than 0.05 was considered for statistical analysis.

Results

The mean age of the participants in this study, was 36.4±13.16 y (range 18-70 y), 112(44.8%) and 138(55.2%)

were male and female respectively. 82.4% were married and 17.6% were single.

The mean BMI of the participants was 25.39±3.33 kg/m² and the mean waist circumference of these individuals was 83.22 ± 12.92 cm, of which 42.8% were overweight, 8.4% were obese and 23.6% had abdominal obesity.

Factor analysis revealed two factors that explained 32.71% of the variance in dietary patterns. The healthy nutritional pattern consisted of poultry, fish, vegetables, organ meat, potatoes, fruits, natural juices, jams and compotes, tomatoes, and dairy, which accounts for 14.74% of the variance in dietary patterns. The unhealthy dietary habit that included sweet snacks, salty snacks, beverages, coffee and tea, mayonnaise, salt, processed meats, refined cereals, red meats, sweets and solid fats accounts for 17.97% of the variance in dietary patterns [Table 2]. Besides, factor loadings of corn, egg, nuts, seeds and pips, liquid fat, and whole grains were less than 0.2 and did not fit into any of the dominant patterns.

Table 2. Dietary patterns and factor load of food patterns derived from principal component analysis

Food groups	dietary patterns	
	Healthy	Unhealthy
Sweet snacks		0.77
Salty snacks		0.75
Soft drinks		0.66
Coffee and tea		0.52
Mayonnaise		0.49
Salt		0.48
Processed meats		0.45
Refined cereals		0.35
Red Meat		0.31
Cookies		0.28
Solid fats		0.21
chicken and fish	0.65	
Vegetables	0.64	
Organ meats	0.63	
Potato	0.49	
Fruits, natural juices	0.46	
Jam and compote	0.39	
Tomato	0.36	
Dairy	0.33	

The KMO index is 0.56; hence, the data are sufficient, and the significance of Bartlett's Spear Test shows there is a significant correlation between food groups [$\chi^2 = 1670.62$ and $p < 0.001$].

As can be seen in the above table, two major dietary patterns were obtained. The most common factor in the

unhealthy nutritional pattern is the sweet snack food group [0.77] and in the healthy dietary pattern, it belongs to the chicken and fish diet [0.65]. Further, the minimum load factor in the unhealthy dietary pattern observes in the solid fat group [0.21] and in the healthy dietary pattern notices in the dairy group [0.33]. The relationship between the healthy and unhealthy eating patterns with the Anthropometric factors of the participants is shown in Table 3.

There were no significant differences between unhealthy diet and height ($p=0.82$) and waist circumference ($p=0.32$) variables. The unhealthy dietary pattern was correlated with body weight ($p=0.003$), hip circumference ($p < 0.001$) and body mass index ($p < 0.001$). Also, there were significant relationship between healthy dietary pattern and height ($p < 0.001$), weight ($p < 0.001$), waist circumference ($p < 0.001$), hip circumference ($p < 0.001$) and body mass index ($p=0.001$).

The unhealthy diet was significantly associated with general obesity ($p < 0.001$) and abdominal obesity ($p=0.01$). Both general and abdominal obesity measures were significantly associated with Healthy diet pattern ($p = 0.003$ and $p < 0.001$ respectively).

Table 3. Anthropometric measures across tertiles of major dietary patterns among Gonabad adults.

Variable	Tertiles of Unhealthy pattern			Tertiles of Healthy pattern		
	Tertile1 (n=83)	Tertile2 (n=83)	Tertile3 (n=84)	Tertile1 (n=83)	Tertile2 (n=84)	Tertile3 (n=83)
Height (cm)	166.60±12.50	167.60±10.51	166.78±9.66	166.69±12.31	171.73±8.65	162.49±9.52
	F=0.20, P=0.82			F=16.93, P<0.001		
Weight (kg)	70.27±14.41	67.51±9.13	73.80±10.96	73.57±14.29	71.46±9.76	66.59±1030
	F=6.06, P=0.003			F=7.88, P<0.001		
Waist (cm)	84.69±13.32	81.97±13.63	84.50±11.71	88.55±13.01	84.66±9.85	77.95±13.46
	F=1.15, P=0.32			F16.03, P<0.001		
Hip (cm)	101.75±10.26	92.20±15.63	97.20±10.01	103.69±9.57	94.79±12.80	92.69±13.06
	F=12.65, P<0.001			F19.92, P<0.001		
BMI (kg/m ²)	25.15±3.10	24.12±3.46	26.85±3.60	26.36±3.44	24.26±3.14	25.66±3.71
	F=11.03, P<0.001			F=7.77, P=0.001		

P-values reported based on ANOVA test. Data described as mean ±SD

Table 4. General and abdominal obesity across tertiles of major dietary patterns among Gonabad adults.

Variable	Groups	N	Tertiles of Unhealthy pattern			Tertiles of Healthy pattern		
			Tertile1 % (n=83)	Tertile2% (n=83)	Tertile3 % (n=84)	Tertile1 % (n=83)	Tertile2% (n=84)	Tertile3% (n=83)
Weight status	Normal	122	23.0	50.8	26.2	22.1	42.7	35.2
	Overweight	107	51.4	15.9	32.7	44.9	26.1	29.0
	Obese	21	0.0	19.0	81.0	38.1	19.0	42.9
	Sig		$\chi^2 = 59.85, P < 0.001$			$\chi^2 = 16.39, P = 0.003$		
Abdominal Obesity	Yes	191	28.3	35.6	36.1	20.4	38.7	40.9
	No	59	49.2	25.4	25.4	74.6	16.9	8.5
	Sig		$\chi^2 = 8.86, P = 0.01$			$\chi^2 = 60.41, P < 0.001$		

P-values reported based on chi-squared test.

Discussion

To our knowledge, this study is the first to identify the dietary patterns among Gonabad people.

In the present study, we report analysis of dietary patterns among 250 adult Gonabad people aged 18–70 and the association of dietary patterns with general and abdominal obesity. Two dietary patterns were identified. The first pattern, the Healthy dietary pattern, was characterized mainly by high consumption of poultry and fish, vegetables, organ meat, potatoes, fruits, natural juices, jams and compotes, tomatoes, and dairy. The second pattern, the Unhealthy dietary pattern, consisted of a combination of sweet snacks, salty snacks, beverages, coffee and tea, mayonnaise, salt, processed meats, refined cereals, red meats, sweets, and solid fats. The identified “Healthy” and “Un healthy” dietary patterns are similar to those reported previously conducted among adults(39-41).

As there were no studies to identify dietary patterns in gonabad, it was not possible to compare the results of this study with other studies. However, the dietary patterns detected in this study are comparable with the findings of other studies in Iran.

Rezazadeh et al. identified two dietary patterns (namely Healthy, and unhealthy) among adult women in Tehran(16). Unhealthy dietary pattern in this study was similar to an Unhealthy dietary pattern with a high intake of refined cereals, processed meats, hydrogenated oil, and sweets.

In Esmaeilzadeh et al, study aimed at identifying the major dietary patterns relation with general obesity and central adiposity among Iranian women. The unhealthy dietary pattern was the same as the unhealthy dietary pattern (with a high intake of refined cereals, processed meats, hydrogenated oil, and sweets) in the present study(42).

The Healthy dietary Pattern in this study is the same as a healthy dietary pattern of Rezazadeh, mirmiran, azadbakht studies, include a variety of fruits, vegetable, and poultry(6, 16, 43).

Despite inconsistencies between the results derived from different studies around the world, there are some similarities regarding dietary patterns identified among the adult people. Sánchez-Villegas A et al. fined two major dietary patterns(44). The Mediterranean diet, with similar components to the present study contained vegetables, fruits, poultry, nuts and potatoes. In Parl mendez et al, study, among American men and women, vegetables, fruits and milk patterns similar to the present study(45).

Also, in McNaughton et al study, dietary patterns of fruits, vegetables and dairy had the same healthy nutritional habits as found in this study(46).

Other studies have indicated dietary patterns can be can varied and depending on the number of the pattern obtained. In Naja et al. study, three dietary patterns were

identified in Lebanese adults: Fast Food/Dessert," "Traditional Lebanese," and "High Protein(47).

The patterns obtained in the present study are derived from the data collected in a particular population of Iran, and therefore, they are not expected to be quite truthful in a population with different eating habits. Because of the geographical, cultural, social and economic differences, these differences are not strange and even the dietary pattern of a city cannot be generalized to the other city of the same country.

In this study, unhealthy dietary pattern was correlated with weight, hip circumference, and body mass index(BMI). Also, individuals in high healthy diet score (tertile 3 group), showed significantly decreased in weight, waist and hip circumference. Individuals in the lowest tertile of the unhealthy dietary pattern score, were less likely to be generally obese whereas, those in the upper tertile of healthy dietary pattern were less likely to be generally and centrally obese.

In the mohammadshahi study, a relationship was established between a healthy diet and a lower body mass index (BMI) and waist circumference. Moreover, the Western dietary pattern has shown a positive correlation with BMI, waist circumference, body fat percentage (27). A healthy diet is usually rich in foods containing low calorie content and more nutrients, so it has a positive effect in weight control. According to a study conducted in Tehran, a healthy dietary pattern had an important role in obesity(48). The results of Safarian and, Mirmiran, showed that adherence to the Western dietary pattern was associated with increased weight, BMI, and waist circumference (47,41). In the study of Rezazadeh et al., it was found that a healthy dietary pattern was inversely associated with BMI and waist circumference(16).

In the study on Iranian women dietary patterns were divided into two healthy and unhealthy pattern and showed that people with the highest quartile of unhealthy dietary patterns were more likely to develop general and abdominal obesity. But people who are in the top quartile range of healthy eating patterns have a lower risk for obesity and abdominal obesity (16, 49).

The results of a cross-sectional study in women teachers in Tehran as well as Mirmiran study showed that adherence to the Western dietary pattern was associated with increased BMI and abdominal obesity. Participants in the highest quartile of a healthy pattern had lower risk of general obesity, also, people with the highest quartiles of unhealthy dietary patterns had higher body mass index (BMI) and general obesity compared to those with the lowest quartiles(43, 50).

The result of aghapour et al showed the participants who had higher scores for the unhealthy and the healthy dietary patterns were more and less likely to have overweight or obesity, respectively, compared to participants with lower scores(24).

There were many similarities between the dietary patterns obtained in our study and other studies regarding the association of dietary patterns with overweight or obesity.

In this study, unhealthy pattern, like the unhealthy pattern of Rezazadeh study(16) and western pattern of study of esmailzadeh (42), was positively associated with obesity risk. However, in the study of Aghapour et al. No association was found between the healthy dietary pattern and weight status in adolescents(24). Dietary patterns, characterized by high fats, sugars and snacks have not been associated with the risk of overweight or obesity in adolescents(51, 52).

Studies in other parts of the world including the United States (53, 54), Europe (45) and Asia (43) also showed an inverse association between healthy dietary pattern and general and abdominal obesity. McNaughton reported, In English women, dietary patterns of fruits, vegetables, and dairy products were inversely correlated with BMI and waist circumference (46). In another study in the United States, Consumption of whole grains has been associated with lower BMI and waist circumference(53).

The inverse association between healthy pattern and risk of general and central obesity was similar to the results reported in the us (55, 56) and European(57) countries. In other studies, the inverse relation between BMI and weight gain and "whole grains, fruits and vegetables" pattern were reported in men(58) and women(59) who participated in the Nurses' Health Study. In addition, a "low-fat dairy products, grains, and fruit" pattern in American women had a reverse association with annual changes in BMI and WC(60, 61). Otherwise, dietary pattern of meat and fat in American men and women(62) and the dietary pattern of meat and pasta in Italian women and men (63) were positively associated with BMI.

The positive relationship between unhealthy dietary pattern and the likelihood of overweight and obesity may be due to the high intake of animal fats and refined cereals in the unhealthy dietary pattern. High fat intake is known as a risk factor for obesity (64) and refined grains with a high glycaemic index can increase hunger and are effective in boosting overeating(65)..

Inverse Relationship between Healthy Pattern Observed in Recent Study with weight, waist circumferences and Abdominal Obesity maybe because of the effect of foods rich in fiber and complex carbohydrates, low glycemic index and low energy density like vegetables, fruits and legumes(66, 67) and low fat consumption on appetite and food intake(68). While, the positive relation between unhealthy pattern weight and general obesity may be due to excessive consumption of higher glycemic index carbohydrates (refined grains, sugars and sweets) which leads to higher glycemic responses and increased fat synthesis and fat accumulation(69). Moreover, these types of carbohydrates increase hunger and stimulate overeating.

On the other hand, increased fat intake, which is observed in unhealthy diets, was considered a risk factor for weight gain. (65).

One of the strengths of the present study was to investigate both sexes and control of the effect of many confounding variables, especially specially dieting and history of diabetes, hypertension and hyperlipidemia.

A number of limitations must be taken into account in the interpretation of our findings. Because of the cross-sectional design of this study, causality cannot be inferred; therefore, our findings need to be confirmed in prospective studies.

Second, the FFQ limits for the assessment of dietary intakes, including the use of memory and the potential for under-reporting and over-reporting, should also be taken into consideration (67).

Conclusion

Two Major dietary Healthy and Unhealthy Dietary patterns were significantly associated with general and abdominal obesity among adults in Gonabad. It is suggested to consider anthropometric measures in determining the relationship between dietary patterns and disease. Further studies with future prospective studies, are required to confirm these findings.

Financial disclosure

The authors declared no financial interest.

Funding/Support

This work was financially supported by the Gonabad University of Medical Sciences, Gonabad, Iran.

References

1. WHO: Obesity and Overweight factsheet from the WHO. Retrieved. December 13, 2018, from <https://www.who.int/news-room/fact-sheets/>. detail/obesity-and-overweight.
2. Djalalinia S, Saeedi Moghaddam S, Sheidaei A, Rezaei N, Naghibi Irvani SS, Modirian M, et al. Patterns of Obesity and Overweight in the Iranian Population: Findings of STEPs 2016. *Frontiers in Endocrinology*. 2020;11:42.
3. Campfield LA, Smith FJ. The pathogenesis of obesity. *Baillieres Best Pract Res Clin Endocrinol Metab*. 1999;13(1):13-30.
4. Reilly JJ, Kelly J. Long-term impact of overweight and obesity in childhood and adolescence on morbidity and premature mortality in adulthood: systematic review. *Int J Obes (Lond)*. 2011;35(7):891-8.
5. Zimmermann-Belsing T, Feldt-Rasmussen U. Obesity: the new worldwide epidemic threat to general health and our complete lack of effective treatment. *Endocrinology*. 2004;145(4):1501-2.
6. Azadbakht L, Esmailzadeh A. Dietary and non-dietary determinants of central adiposity among Tehrani women. *Public Health Nutr*. 2008;11(5):528-34.

7. Azadbakht L, Mirmiran P, Shiva N, Azizi F. General obesity and central adiposity in a representative sample of Tehranian adults: prevalence and determinants. *Int J Vitam Nutr Res*. 2005;75(4):297-304.
8. Hoffmann K, Schulze M, Boeing H, Altenburg H. Dietary patterns: report of an international workshop. *Public health nutrition*. 2002;5(1):89-90.
9. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Current opinion in lipidology*. 2002;13(1):3-9.
10. Newby PK, Tucker KL. Empirically derived eating patterns using factor or cluster analysis: a review. *Nutr Rev*. 2004;62(5):177-203.
11. Hu FB. Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol*. 2002;13(1):3-9.
12. Maskarinec G, Novotny R, Tasaki K. Dietary patterns are associated with body mass index in multiethnic women. *The Journal of nutrition*. 2000;130(12):3068-72.
13. Motswagole B, Jackson J, Kobue-Lekalake R, Maruapula S, Mongwaketse T, Kwape L, et al. The Association of General and Central Obesity with Dietary Patterns and Socioeconomic Status in Adult Women in Botswana. *Journal of Obesity*. 2020;2020.
14. Newby P, Muller D, Hallfrisch J, Andres R, Tucker KL. Food patterns measured by factor analysis and anthropometric changes in adults. *The American journal of clinical nutrition*. 2004;80(2):504-13.
15. Newby PK, Muller D, Hallfrisch J, Qiao N, Andres R, Tucker KL. Dietary patterns and changes in body mass index and waist circumference in adults. *The American journal of clinical nutrition*. 2003;77(6):1417-25.
16. Rezazadeh A, Rashidkhani B. The association of general and central obesity with major dietary patterns of adult women living in Tehran, Iran. *J Nutr Sci Vitaminol (Tokyo)*. 2010;56(2):132-8.
17. van Dam RM, Grievink L, Ocké MC, Feskens EJ. Patterns of food consumption and risk factors for cardiovascular disease in the general Dutch population. *The American journal of clinical nutrition*. 2003;77(5):1156-63.
18. Yu C, Shi Z, Lv J, Du H, Qi L, Guo Y, et al. Major dietary patterns in relation to general and central obesity among Chinese adults. *Nutrients*. 2015;7(7):5834-49.
19. Vaisi-Raygani A, Mohammadi M, Jalali R, Ghobadi A, Salari N. The prevalence of obesity in older adults in Iran: a systematic review and meta-analysis. *BMC geriatrics*. 2019;19(1):1-9.
20. Willett W. *Nutritional epidemiology*: Oxford university press; 2012.
21. Annis NM, Cash TF, Hrabosky JI. Body image and psychosocial differences among stable average weight, currently overweight, and formerly overweight women: the role of stigmatizing experiences. *Body Image*. 2004;1(2):155-67.
22. Thorpe KE, Florence CS, Howard DH, Joski P. The impact of obesity on rising medical spending. *Health Aff (Millwood)*. 2004;Suppl Web Exclusives:W4-480-6.
23. Gokee-LaRose J, Gorin AA, Raynor HA, Laska MN, Jeffery RW, Levy RL, et al. Are standard behavioral weight loss programs effective for young adults? *International Journal of Obesity*. 2009;33(12):1374-80.
24. Aghapour B, Rashidi A, Dorosti-Motlagh A, Mehrabi Y. The association between major dietary patterns and overweight or obesity among Iranian adolescent girls. *Iranian Journal of Nutrition Sciences & Food Technology*. 2013;7(5):289-99.
25. Freeland-Graves J, Nitzke S. Position of the American Dietetic Association: total diet approach to communicating food and nutrition information. *J Am Diet Assoc*. 2002;102(1):100-8.
26. Tarighat-Esfanjani A, Ebrahimi-Mameghani M, Asghari-Jafarabadi M, Jafari-Vayghan H, Ghadimi SS. Identification of major dietary patterns in Tabriz adults. *Medical Journal of Tabriz University of Medical Sciences*. 2016 Jun 12;38(2):48-55.
27. Mohammad Shahi M, Heidari F, Mola K, Helli B, Ijadi M, Amirian Z, et al. Association of Dietary Patterns and Indicators of Disease Activity in Patients with Rheumatoid Arthritis. *Iranian Journal of Nutrition Sciences & Food Technology*. 2014;9(3):9-20.
28. Dixon JK. *Statistical methods for health care research*: Munro BH, editor., *Exploratory Factor Analysis*. 5th ed. Philadelphia: Lippincott Williams & Wilkins, 2005:321-50.
29. Moshkani RF, Saneei P, Esmailzadeh A, Keshteli AH, Feizi A, Adibi P. Association between patterns of dietary habits and obesity in Iranian adults. *Iranian Journal of Nutrition Sciences & Food Technology*. 2016;11(2):19-34.
30. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public health nutrition*. 2010;13(5):654-62.
31. Zhang C-X, Ho SC, Fu J-H, Cheng S-Z, Chen Y-M, Lin F-Y. Dietary patterns and breast cancer risk among Chinese women. *Cancer causes & control*. 2011;22(1):115-24.
32. Hosseini Esfahani F, Jazayeri A, Mirmiran P, Mehrabi Y, Azizi F. Dietary patterns and their association with socio-demographic and lifestyle factors among Thehrani adults: Tehran Lipid and Glucose Study. *Journal of School of Public Health and Institute of Public Health Research*. 2008;6(1):23-36.
33. Organization WH. *Waist circumference and waist-hip ratio: report of a WHO expert consultation*, Geneva, 8-11 December 2008. 2011.
34. Babai MA, Arasteh P, Hadibarhaghtalab M, Naghizadeh MM, Salehi A, Askari A, et al. Defining a BMI cut-off point for the Iranian population: the Shiraz Heart Study. *PloS one*. 2016;11(8):e0160639.
35. Chang Y, Ryu S, Suh B, Yun K, Kim C, Cho S. Impact of BMI on the incidence of metabolic abnormalities in metabolically healthy men. *International journal of obesity*. 2012;36(9):1187-94.
36. Esteghamati A, Abbasi M, Rashidi A, Meysamie A, Khalilzadeh O, Haghazali M, et al. Optimal waist circumference cut-offs for the diagnosis of metabolic syndrome in Iranian adults: results of the third national survey of risk factors of non-communicable diseases (SuRFNCD-2007). *Diabetic Medicine*. 2009;26(7):745-6.
37. WHO EC. *Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies*. *Lancet (London, England)*. 2004;363(9403):157.
38. Zandieh A, Esteghamati A, Morteza A, Noshad S, Khalilzadeh O, Gouya MM, et al. Appropriate BMI cut-off values for identification of metabolic risk factors: third national surveillance of risk factors of non-communicable diseases in Iran (SuRFNCD-2007). *Annals of human biology*. 2012;39(6):484-9.

39. Newby PK, Weismayer C, Akesson A, Tucker KL, Wolk A. Longitudinal changes in food patterns predict changes in weight and body mass index and the effects are greatest in obese women. *J Nutr.* 2006;136(10):2580-7.
40. Gholami A, Sohrabi M, Abbasi-Ghahramanloo A, Moradpour F, Safiri S, Maadi M, et al. Identifying the pattern of unhealthy dietary habits among an Iranian population: A latent class analysis. *Medical journal of the Islamic Republic of Iran.* 2018;32:69.
41. Rezazadeh A, Hassanpour R, Rashidkhani B. Major Dietary Patterns of Women Living in Urban and Rural Areas of Dashtestan County (Bushehr Province), 2017. *Iranian South Medical Journal.* 2020;23(2):129-42.
42. Esmailzadeh A, Azadbakht L. Major dietary patterns in relation to general obesity and central adiposity among Iranian women. *J Nutr.* 2008;138(2):358-63.
43. mirmiran P, Djazayeri A, Hosseini esfahani F, Mehrabi Y, Azizi F. Change in food patterns of Tehrani adults and its association with changes in their body weight and body mass index in District 13 of Tehran: Tehran Lipid and Glucose Study. *Iranian Journal of Nutrition Sciences & Food Technology.* 2008;2(4):67-80.
44. Sánchez-Villegas A, Delgado-Rodríguez M, Martínez-González MÁ, de Irala-Estévez J, for the SUNg. Gender, age, socio-demographic and lifestyle factors associated with major dietary patterns in the Spanish Project SUN (Seguimiento Universidad de Navarra). *European Journal of Clinical Nutrition.* 2003;57(2):285-92.
45. Mendez MA, Popkin BM, Jakszyn P, Berenguer A, Tormo MJ, Sanchez MJ, et al. Adherence to a Mediterranean diet is associated with reduced 3-year incidence of obesity. *J Nutr.* 2006;136(11):2934-8.
46. McNaughton SA, Mishra GD, Stephen AM, Wadsworth ME. Dietary patterns throughout adult life are associated with body mass index, waist circumference, blood pressure, and red cell folate. *J Nutr.* 2007;137(1):99-105.
47. Naja F, Nasreddine L, Itani L, Adra N, Sibai AM, Hwalla N. Association between dietary patterns and the risk of metabolic syndrome among Lebanese adults. *Eur J Nutr.* 2013;52(1):97-105.
48. Azadbakht L, Mirmiran P, Hosseini F, Azizi F. Diet quality status of most Tehranian adults needs improvement. *Asia Pac J Clin Nutr.* 2005;14(2):163-8.
49. Rezazadeh A, Rashidkhani B, Omidvar N. Investigation of predominant dietary patterns, general obesity and central obesity in adults living in north of Tehran. *J Shahid Beheshti Univ Med Sci.* 2009;4:246-57. [in Persian].
50. Hosseini Esfahani F, Mirmiran P, Djazayeri S, Mehrabi Y, Azizi F. Change in Food Patterns and its Relation to Alterations in Central Adiposity in Tehranian of District 13 Adults. *Iranian Journal of Endocrinology and Metabolism.* 2008;10(4):299-312.
51. Craig LC, McNeill G, Macdiarmid JI, Masson LF, Holmes BA. Dietary patterns of school-age children in Scotland: association with socio-economic indicators, physical activity and obesity. *Br J Nutr.* 2010;103(3):319-34.
52. McNaughton SA, Ball K, Mishra GD, Crawford DA. Dietary Patterns of Adolescents and Risk of Obesity and Hypertension. *The Journal of Nutrition.* 2008;138(2):364-70.
53. Khani BR, Ye W, Terry P, Wolk A. Reproducibility and validity of major dietary patterns among Swedish women assessed with a food-frequency questionnaire. *J Nutr.* 2004;134(6):1541-5.
54. Sanchez-Villegas A, Delgado-Rodriguez M, Martinez-Gonzalez MA, De Irala-Estevéz J. Gender, age, socio-demographic and lifestyle factors associated with major dietary patterns in the Spanish Project SUN (Seguimiento Universidad de Navarra). *Eur J Clin Nutr.* 2003;57(2):285-92.
55. Lohman TG, Roche AF, Martorell R. Anthropometric standardization reference manual. 1st ed. Champaign, IL: Human Kinetics Books; 1988. p. 28-80.
56. Aadahl M, Jørgensen T. Validation of a new self-report instrument for measuring physical activity. *Med Sci Sports Exerc.* 2003;35(7):1196-202.
57. Okubo H, Sasaki S, Murakami K, Kim MK, Takahashi Y, Hosoi Y, et al. Three major dietary patterns are all independently related to the risk of obesity among 3760 Japanese women aged 18-20 years. *Int J Obes (Lond).* 2008;32(3):541-9.
58. Hu FB, Rimm EB, Stampfer MJ, Ascherio A, Spiegelman D, Willett WC. Prospective study of major dietary patterns and risk of coronary heart disease in men. *Am J Clin Nutr.* 2000;72(4):912-21.
59. Schulze MB, Fung TT, Manson JE, Willett WC, Hu FB. Dietary patterns and changes in body weight in women. *Obesity (Silver Spring).* 2006;14(8):1444-53.
60. Newby PK, Muller D, Hallfrisch J, Qiao N, Andres R, Tucker KL. Dietary patterns and changes in body mass index and waist circumference in adults. *Am J Clin Nutr.* 2003;77(6):1417-25.
61. Yang EJ, Kerver JM, Song WO. Dietary patterns of Korean Americans described by factor analysis. *J Am Coll Nutr.* 2005;24(2):115-21.
62. Park SY, Murphy SP, Wilkens LR, Yamamoto JF, Sharma S, Hankin JH, et al. Dietary patterns using the Food Guide Pyramid groups are associated with sociodemographic and lifestyle factors: the multiethnic cohort study. *J Nutr.* 2005;135(4):843-9.
63. Pala V, Sieri S, Masala G, Palli D, Panico S, Vineis P, et al. Associations between dietary pattern and lifestyle, anthropometry and other health indicators in the elderly participants of the EPIC-Italy cohort. *Nutrition, metabolism, and cardiovascular diseases : NMCD.* 2006;16(3):186-201.
64. Lissner L, Heitmann BL. Dietary fat and obesity: evidence from epidemiology. *Eur J Clin Nutr.* 1995;49(2):79-90.
65. Roberts SB. High-glycemic index foods, hunger, and obesity: is there a connection? *Nutr Rev.* 2000;58(6):163-9.
66. Bell EA, Castellanos VH, Pelkman CL, Thorwart ML, Rolls BJ. Energy density of foods affects energy intake in normal-weight women. *Am J Clin Nutr.* 1998;67(3):412-20.
67. Jenkins DJ, Kendall CW, Augustin LS, Franceschi S, Hamidi M, Marchie A, et al. Glycemic index: overview of implications in health and disease. *Am J Clin Nutr.* 2002;76(1):266s-73s.
68. Kimm SY. The role of dietary fiber in the development and treatment of childhood obesity. *Pediatrics.* 1995;96(5 Pt 2):1010-4.
69. Sichieri R. Dietary patterns and their associations with obesity in the Brazilian city of Rio de Janeiro. *Obes Res.* 2002;10(1):42-8.