Nutrition and Food in Health and Disease Vol 12, No 4, Oct-Dec 2025, pages:33-39

*Original Article*

# The Effect of Short-term Green Tea Supplementation on Serum Cortisol and Serotonin Levels in Young Anxious Women

Fatemeh Dini[1], Hassan Pourrazi[2*]

1- M.Sc. in Exercise Physiology, Department of Sport Sciences, Faculty of Social Sciences, Imam Khomeini International University, Qazvin, Iran.
2- Associate Professor of Exercise Physiology, Department of Sport Sciences, Faculty of Social Sciences, Imam Khomeini International University, Qazvin, Iran.

**Received:** March 2025 **Accepted:** April 2025

## ABSTRACT

**Background and Objectives:** In recent years, the use of dietary supplements to combat anxiety has increased. Therefore, the present study aimed to assess the effect of short-term green tea supplementation on serum cortisol and serotonin levels in young anxious women.

**Materials and Methods:** In this randomized clinical trial, 20 young anxious women were randomly assigned to either a control or a green-tea supplementation group. Participants in the supplementation group consumed 1000 mg of green tea daily for one week. Heart rate, systolic blood pressure, rate-pressure product, serum cortisol and serotonin levels were assessed before and after the intervention. Data were analyzed using paired and independent t-tests at a significance level of 0.05.

**Results:** The findings indicated no significant differences between the control and supplementation groups for changes in heart rate, systolic blood pressure and cortisol levels ($P = 0.09$, $P = 0.15$ and $P = 0.96$, respectively). However, green tea supplementation significantly decreased the rate-pressure product and significantly increased serum serotonin levels ($P = 0.05$, $P < 0.0001$, respectively).

**Conclusions:** Overall, short-term green tea supplementation seemed to contribute to anxiety management in young anxious women by decreasing the rate-pressure product and increasing serum serotonin levels.

**Keywords:** Green tea, Cortisol, Serotonin, Anxiety

## Highlights

- Short-term daily consumption of 1000 mg of green tea resulted in a significant increase in serum serotonin levels in young anxious women.
- Green tea supplementation seemed to contribute to anxiety management in young anxious women by decreasing the rate-pressure product.
- Short-term consumption of green tea did not have a significant effect on cortisol levels. However, the downward trend in cortisol concentrations suggests a potential role for green tea in stress modulation.

## Introduction

Anxiety is a psychological condition characterized by a state of anticipatory fear (1). Based on the latest estimates, approximately 60 million individuals in Europe suffer from anxiety disorders. Epidemiological studies indicate that in the United States, the lifetime prevalence of any anxiety disorder is approximately one-third, surpassing that of any other diagnostic category. Anxiety is a significant risk factor for diminished quality of life, increased

***Address for correspondence:** Hassan Pourrazi, Associate Professor of Exercise Physiology, Department of Sport Sciences, Faculty of Social Sciences, Imam Khomeini International University, Qazvin, Iran. Email Address: pourrazi@soc.ikiu.ac.ir

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

mortality rates and various physical health complications, particularly cardiovascular diseases (CVD). Survey data from the United States suggest that women are twice as likely as men to develop panic disorder or phobias throughout their lifetime. Although gender-based differences in anxiety disorders are relatively subtle, evidence indicates a increasing prevalence of anxiety in women. The exacerbation of anxiety symptoms, whether emotional (e.g., fear and restlessness) or physiological (e.g., palpitations and tremors), represents a common hallmark within all anxiety disorders (1,2). Physiological perspective on anxiety from a physiological standpoint, research indicates that anxiety activates the sympathetic nervous system, affecting several physiological markers such as heart rate and blood pressure, leading to an increase in the two parameters (3). This phenomenon may increase myocardial oxygen demand, with the rate-pressure product (RPP) as one of its key indicators (4,5). Additionally, the assessment of other physiological markers, including specific hormones such as cortisol, serotonin and endorphins, serves as an effective approach for diagnosing anxiety-linked disorders (2). From these, cortisol is a hormone that regulates metabolism, preparing organs to deal with various conditions by enhancing the activity of other stress-linked hormones such as adrenaline and noradrenaline. Cortisol is widely documented as a key biomarker of stress and due to high levels of stress, the human body is frequently exposed to increased cortisol levels (2,6). On the contrary, neurotransmitters and mediators such as serotonin and endorphins are regarded as biomarkers of well-being and happiness, playing a crucial role in emotional regulation and mental health. Serotonin (5-hydroxytryptamine) was first discovered in 1948 in blood platelets and its presence in the brain was identified in later years. Then, extensive research has been carried out on its role in psychiatric disorders. Serotonin is associated with several psychological functions, one of which is anxiety-linked disorders (7). Based on the high prevalence of anxiety in today's society, particularly in women, researchers continuously search for effective strategies for its prevention and management. In recent years, the effect of various dietary and herbal supplements on anxiety has received significant attention in the health research field. From these, green tea supplementation has emerged as a widely used herbal remedy for promoting relaxation and improving mental well-being. Recent studies have investigated the effects of green tea on mental disorders, including anxiety and stress. Conger et al. (2019) reported that even a small intake of green tea could include a positive psychological effect. For example, drinking hot green tea has been shown to enhance mood (8). Green tea is derived from the leaves and shoots of the *Camellia sinensis* plant and contains high quantities of catechins, minerals and various vitamins. Catechins are potent antioxidants, while flavonoids, theanine and aromatic compounds are key constituents of green tea. Recent studies indicate that theanine, found in green tea, can decrease stress by inhibiting cortical nerve excitation in humans or by affecting neurotransmitter secretion and function in the central nervous system (CNS) (9-12). Epidemiological studies have indicated a connection between green tea consumption and a decrease in psychological distress. Research on the nutrients in green tea, especially flavonoids such as epigallocatechin gallate (EGCG) and L-theanine, has yielded varied results. Components of green tea modulate acute brain oscillatory activity. The L-theanine, present in green tea, includes anxiolytic effects, increasing alpha brain-wave activity in humans and preventing an increase in blood pressure. The EGCG, another constituent of green tea, enhances alpha and beta wave fluctuations. However, in another study, glutamate (Glu) of green tea included no effect on stress in rats (13-17). Based on the contradictory results regarding the effects of green tea on anxiety and increasing prevalence of anxiety, particularly in women, further research is critical. Based on existing reviews, only a few studies have investigated effects of green tea supplementation on anxiety-linked hormonal markers in women, with a very limited data available in this area. Therefore, the present study aimed to investigate effects of short-term green tea supplementation on serum cortisol and serotonin levels, as well as other physiological markers, in young anxious women.

## Materials and Methods

The present study was a randomized clinical trial carried out in a pretest-posttest format within a control group. The study population consisted of young anxious women aged 18–22 y. Participants were screened and

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

selected using Spielberger anxiety inventory, which included questions on trait and state anxieties. Based on the anxiety scores achieved as the primary inclusion criteria and considering the nature and objectives of the study, 20 young anxious women were voluntarily selected and randomly assigned to two groups of A) control group (ten participants) and B) green-tea supplementation group (ten participants). All procedures of the study were carried out in accordance with the ethical principles of the Medical Ethics Committee, Qazvin University of Medical Sciences, Qazvin, Iran (IR.QUMS.REC.1403.031).

The participants in the supplementation group consumed the specified quantities of green tea supplementation daily. The supplement consisted of green-tea herbal tablets, which were purchased in bulk from Dineh, Iran. Each tablet contained 500 mg of green-tea leaf powder and 50 mg of total polyphenols. Participants in the supplementation group consumed two tablets daily for 1 w, resulting in a daily green tea dose of 1000 mg (18). The timing and manner of supplement consumption were carefully controlled by the researcher.

Before and after the supplementation time, several physical and physiological indicators of the participants, including height, weight, body mass index (BMI), heart rate, systolic and diastolic blood pressures, rate-pressure product and cortisol and serum serotonin levels, were assessed. Additionally, the menstrual cycle of the participants was carefully controlled through self-report, with the assessments during the luteal phase (nearly Days 17–25). To achieve blood pressure and heart rate values, participants were asked to visit the medical venue at the scheduled time. Heart rate and blood pressure were assessed using digital upper-arm blood pressure monitor (Beurer BM96, Germany). The procedure was as follows: the participant sat in a completely relaxed position on a chair for 10-15 min before the assessment. The device was then attached to the left wrist based on standard procedures, with the arm positioned at heart level. The device was activated and the heart rate and systolic and diastolic blood pressure were recorded. The myocardial oxygen demand was calculated by multiplying the systolic blood pressure by the heart rate.

Then, 3 ml of blood (without anticoagulant) was collected from the antecubital vein (orange) using 5-ml syringes. The sample was then set at room temperature (RT) (22–25°C) for 30 min to clot. Then, the serum was separated using centrifuge and cortisol levels were assessed using Architect cortisol kit E-Abbott 8D15, Germany, while serotonin levels were assessed using ELISA kit, Zellbio, Germany. The two hormone levels were quantified using auto-analyzer and reported in international units (IU) per liter. After data collection, normality of data distribution was assessed using Shapiro-Wilk test. Data were analyzed using paired and independent t-tests. Statistical analyses were carried out using SPSS v.22 (IBM, USA), with a significance level set at 0.05 ($P < 0.05$).

## Results

Table 1 presents the participants' demographic characteristics for mean and standard deviation (SD). The collected data, based on the Shapiro-Wilk test, showed a normal distribution ($P < 0.05$).

Results of the paired t-test indicated no significant differences between the pre and post-test values for heart rate, systolic blood pressure, rate-pressure product and serum cortisol in the control and green-tea supplementation groups (Table 2). However, a significant difference was observed between the pre and post-test values for serum serotonin in the green-tea supplementation group, with serum serotonin levels significantly increased in this group (Table 2). Furthermore, the findings showed no significant differences between the control and green-tea supplementation groups for the range of changes in heart rate, systolic blood pressure and cortisol (Table 3). However, green tea supplementation led to a significant decrease in the range of changes in rate-pressure product and a significant increase in the range of changes in serum serotonin (Table 3).

**Table 1**. Participants' demographic characteristics (mean ±SD) by group

| Variable | Control Group | Supplement Group | P |
|---|---|---|---|
| Age (years) | 20.6 ± 1.89 | 20.3 ± 1.25 | 0.99 |
| Height (cm) | 162.1 ± 7.2 | 160.9 ± 7.54 | 0.98 |
| Body Weight (kg) | 59.75 ± 12.54 | 62.25 ± 13.86 | 0.67 |
| Body Mass Index (BMI) | 23.04 ± 4.15 | 24.6 ± 5.19 | 0.71 |
| Body Fat Percentage (%) | 27.44 ± 5.77 | 28.43 ± 6.07 | 0.86 |
| Anxiety Score | 52.70 ± 8.9 | 54.75 ± 5.58 | 0.73 |

SD, Standard deviation

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

**Table 2**. Variables assessed in the control and green-tea supplement groups during pre and post-tests

| Variable | Group | Stage | Δ (pre-post) | t | P |
|---|---|---|---|---|---|
| Heart Rate (beats/min) | Control | Pre-test | 78.5 ± 2.8 | -0.19 | 0.84 |
| | | Post-test | 78.6 ± 9.7 | | |
| | Supplement | Pre-test | 82.6 ± 6.02 | 1.7 | 0.86 |
| | | Post-test | 80.9 ± 9.5 | | |
| Systolic Blood Pressure (mmHg) | Control | Pre-test | 111.01 ± 9.6 | -1.96 | 0.081 |
| | | Post-test | 114.3 ± 6 | | |
| | Supplement | Pre-test | 111.4 ± 9.3 | 0.6 | 0.77 |
| | | Post-test | 110.5 ± 8.5 | | |
| Double Product (Heart Rate * Systolic Pressure) | Control | Pre-test | 8733.9 ± 1297.8 | -1.82 | 0.10 |
| | | Post-test | 8996.1 ± 1124.4 | | |
| | Supplement | Pre-test | 9222.8 ± 1328.4 | 1.24 | 0.24 |
| | | Post-test | 8979.4 ± 988.9 | | |
| Serum Cortisol (µg/dL) | Control | Pre-test | 8.5 ± 3.09 | 0.33 | 0.74 |
| | | Post-test | 8.4 ± 2.7 | | |
| | Supplement | Pre-test | 9.27 ± 2.77 | 0.22 | 0.82 |
| | | Post-test | 9.14 ± 2.78 | | |
| Serum Serotonin (ng/mL) | Control | Pre-test | 139.4 ± 4.58 | 1.11 | 0.29 |
| | | Post-test | 138.3 ± 3.1 | | |
| | Supplement | Pre-test | 132.6 ± 18.43 | -12.4 | 0.07* |
| | | Post-test | 145.89 ± 21.89 | | |

*Significant pre- and post-test difference in the supplement group (P < 0.01)

**Table 3**. Changing ranges of variables in the control and green-tea supplement groups

| Variable | Mean difference | t | P |
|---|---|---|---|
| Heart Rate (beats/min) | 1.8 ± 0.2 | 1.77 | 0.13 |
| Systolic Blood Pressure (mmHg) | 3.9 ± 0.6 | 1.49 | 0.15 |
| Double Product (Heart Rate * Systolic Pressure)# | 506.9 ± 87.2 | 6.14 | 0.05 |
| Serum Cortisol (µg/dL) | 0.03 ± 0.06 | 0.04 | 0.96 |
| Serum Serotonin (ng/mL)# | -13.5 ± 1.15 | -11.7 | 0.0001 |

#Significant difference with the control group (P < 0.01)

## Discussion

Anxiety is a common issue in modern life that can include negative effects on individuals' quality of life. One of the natural and effective methods to decrease anxiety is the use of natural dietary supplements. Therefore, the present study was carried out to investigate the effects of green tea supplementation on serum cortisol and serotonin levels, as well as physiological markers, in young anxious women. The results of this study indicated that daily consumption of 1000 mg of green tea for 1 w significantly decreased the range of changes in the double product and significantly increased the range of changes in serum serotonin, compared to that consumption of a placebo did. Although no significant changes were observed in heart rate, systolic blood pressure and serum cortisol in young anxious women following green tea consumption, these markers decreased after 1 w of supplementation with green tea. In a study by Wu et al. (2022), the effect of green tea

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

on the prevention of high blood pressure in mice on high-salt diets was assessed. The findings of this study showed that green tea supplements played an important role in preventing the increase in blood pressure caused by a high-salt diet. Particularly, these supplements activate the phosphoinositide 3-kinase/protein kinase B signaling pathway, which modulates the gene expression linked to blood pressure regulation. Additionally, green tea supplementation showed significant potentials in improving tissue damages in vital organs such as heart, liver and kidneys (19). Similarly, Ta'ati and colleagues (2021) reported that 3 w of green tea consumption resulted in a decrease in blood pressure and heart rate in sedentary middle-aged women. Similar to the present study, Nogueira et al. (2017) reported that daily consumption of 500 mg of green tea supplementation for 1 w did not result in significant differences in systolic blood pressure; however, the consumption for 4 w led to significant decreases (20,21).

Green tea is rich in antioxidant polyphenols such as catechins and flavonols and green tea extract includes a vasodilatory effect, the two of which can be beneficial for cardiovascular health. These physiological effects of green tea can affect CVDa, including hypertension, and may lead to a decrease in systolic blood pressure by up to 5 mm Hg. In animal studies, green tea supplementation and EGCG, the major catechin in green tea, have been reported to prevent increases in blood pressure. To enhance the effectiveness of green tea supplementation and its extract on blood pressure and provide strong recommendations for adults, high-quality, randomized controlled trials with large sample sizes are essential (22).

The present study showed that the consumption of 1000 mg of green tea per day for 1 w did not include a significant effect on serum cortisol levels, although a decrease in cortisol was further pronounced in the supplement group. These findings differed from those of other studies that have reported a positive effect of green tea on decreasing cortisol levels. Almohdi et al. (2022) reported that the consumption of six cups of green tea for 6 w resulted in a significant decrease in cortisol levels and stress (23).

Researchers believe that the catechins and polyphenols in green tea, especially in the decaffeinated variety, play a key role in decreasing cortisol levels and improving stress. These antioxidant and anti-inflammatory compounds help regulate the nervous system's activity and decrease inflammatory responses in the body (24). Studies suggest that EGCG can inhibit the activity of 11,β-hydroxysteroid dehydrogenase type 1 (11,β-HSD1) at the cellular level; thereby, stopping the conversion of cortisone to cortisol. By inhibiting this enzyme, EGCG enables green tea to indirectly decrease the active cortisol level (25). A study carried out by Silva et al. (2025) demonstrated that L-theanine can affect responses to acute stress. This research assessed its effects on tissue cortisol levels, oxidative stress markers and behavior in zebrafish. The findings indicated that L-theanine at a concentration of 45 mg $l^{-1}$, prevented the increase of cortisol levels (26). However, no significant effect on cortisol levels was observed in this study, which might be attributed to the short duration of consumption and the lower dose of green tea, compared to previous studies. Additionally, individual differences in response to the active compounds in green tea could be a factor contributing to these varying results.

In contrast to cortisol, results of the present study indicated that green tea consumption significantly increased serum serotonin levels in young anxious women. Similar to the current study, Ikram and colleagues (2017) investigated the effects of green tea on behavior and neurochemical profiles in rats exposed to chronic stress. The results showed that although food intake decreased in the two groups of rats (green-tea and water groups), the anxiety-linked effects of stress decreased in the rats that consumed green tea. Additionally, serotonin levels in these animals shifted to normal. Overall, this study suggests that the beneficial effects of green tea in managing stress are linked to changes in serotonin metabolism (27). Moreover, a study carried out by Mirza et al. (2013) demonstrated that 5 w of green tea consumption led to an increase in serotonin and a decrease in anxiety symptoms in rats (28). Another study investigating the acute effects of green tea extract (GTE) showed that even a single dose could increase serotonin levels in plasma. These findings, focusing on the short-term effects of green tea consumption, indicated its potential ability to regulate various metabolite levels (29). The L-theanine, an active compound in green tea, plays a significant role in decreasing stress by relieving symptoms of depression and stress linked to psychosocial conditions, improving stress hormone imbalances such as cortisol, decreasing caffeine effects on serotonin and regulating serotonin and glutamate system, which suppresses excessive glutamate excitation, hence improving brain function (24,30). Additionally, a study showed that EGCG included significant effects on serotonin levels. In the group treated with EGCG, serotonin levels in the hippocampus (a brain region linked to mood) increased, while serotonin levels in the gut decreased. These changes suggested that EGCG might help regulate serotonin levels in the brain and gut, exerting antidepressant effects through this neurotransmitter regulation (31). In a study by Seo et al. (2017), which assessed fermented green tea (FGT) consumption, increases in serotonin and improved energy metabolism were observed (32). However, findings of some previous studies were not similar to the findings of the current study. For example, studies by Elseweidy et al. (2009) and Zhang et al. (2021) showed that consuming green tea extract for 10 w and green tea for 12 w resulted in decreased serotonin levels (33,34). These differences might

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

be due to variations in the concentration of active compounds or insufficient bioavailability of these compounds in the supplement or they might be resulted from individual differences. Moreover, the lack of significant within-group and between-group differences in certain variables might be attributed to the relatively small sample size of the present study. However, definitive conclusions need further precise studies.

**Conclusion**

The results of this study indicated that the daily supplementation of 1000 mg of green tea for 1 w led to a significant increase in serum serotonin levels in young women with anxiety, whereas no statistically significant changes were observed in cortisol levels. Nevertheless, a downward trend in cortisol concentrations was recorded in the supplementation group, suggesting a potential role of green tea in stress modulation. Due to its high content of bioactive compounds including catechins and polyphenols, green tea might include beneficial effects on cardiovascular health. Similar to other studies, findings of this investigation underscored the necessity of large-scale, long-term randomized controlled trials to clarify the underlying physiological mechanisms of green tea effects and establish optimal dose and duration parameters for its consumption.

## Financial disclosure

The authors declared no financial interest.

## Funding/Suppor

This study did not receive any grants from any organization.

## References

1. Stonerock GL, Hoffman BM, Smith PJ, Blumenthal JA. Exercise as treatment for anxiety: systematic review and analysis. Annals of behavioral medicine. 2015 Aug 1;49(4):542-56.
2. Łoś K, Waszkiewicz N. Biological markers in anxiety disorders. Journal of Clinical Medicine. 2021 Apr 17;10(8):1744.
3. Roos AL, Goetz T, Krannich M, Donker M, Bieleke M, Caltabiano A, Mainhard T. Control, anxiety and test performance: Self- reported and physiological indicators of anxiety as mediators. British Journal of Educational Psychology. 2023 Apr;93:72-89.
4. Herlekar SS. Revisiting Cardiovascular Reflexes and Resetting in Healthy Adolescents to Analyse Mechanisms of Physiological and Pathophysiological Factors Affecting Individual Blood-Pressure Components. Journal of Stress Physiology & Biochemistry. 2021;17(2):45-54.
5. Vermeulen TD, Boulet LM, Stembridge M, Williams AM, Anholm JD, Subedi P, Gasho C, Ainslie PN, Feigl EO, Foster GE. Influence of myocardial oxygen demand on the coronary vascular response to arterial blood gas changes in humans. American Journal of Physiology-Heart and Circulatory Physiology. 2018 Jul 1;315(1):H132-40.
6. Narmandakh A, Roest AM, de Jonge P, Oldehinkel AJ. Psychosocial and biological risk factors of anxiety disorders in adolescents: a TRAILS report. European child & adolescent psychiatry. 2021 Dec;30:1969-82.
7. Kahn RS, Van Praag HM, Wetzler S, Asnis GM, Barr G. Serotonin and anxiety revisited. Biological Psychiatry. 1988 Jan 15;23(2):189-208.
8. Conger JZ, Singg S. Effects of green tea consumption on psychological health. Therapeutic Advances in Cardiology. 2019;2(2):251-5.
9. Naveed M, BiBi J, Kamboh AA, Suheryani I, Kakar I, Fazlani SA, FangFang X, Yunjuan L, Kakar MU, Abd El-Hack ME, Noreldin AE. Pharmacological values and therapeutic properties of black tea (Camellia sinensis): A comprehensive overview. Biomedicine & Pharmacotherapy. 2018 Apr 1;100:521-31.
10. Ahammed GJ, Li X. Hormonal regulation of health-promoting compounds in tea (Camellia sinensis L.). Plant Physiology and Biochemistry. 2022 Aug 15;185:390-400.
11. Xie X, Wan J, Zheng X, Pan W, Yuan J, Hu B, Feng M, Liu Z, Cai S. Synergistic effects of epigallocatechin gallate and l-theanine in nerve repair and regeneration by anti-amyloid damage, promoting metabolism and nourishing nerve cells. Frontiers in Nutrition. 2022 Aug 10;9:951415.
12. Unno K, Nakamura Y. Green tea suppresses brain aging. Molecules. 2021 Aug 12;26(16):4897.
13. Boyle NB, Billington J, Lawton C, Quadt F, Dye L. A combination of green tea, rhodiola, magnesium and B vitamins modulates brain activity and protects against the effects of induced social stress in healthy volunteers. Nutritional neuroscience. 2022 Sep 2;25(9):1845-59.
14. Yoto A, Motoki M, Murao S, Yokogoshi H. Effects of L-theanine or caffeine intake on changes in blood pressure under physical and psychological stresses. Journal of physiological anthropology. 2012 Dec;31(1):1-9.
15. Unno K, Furushima D, Tanaka Y, Tominaga T, Nakamura H, Yamada H, Taguchi K, Goda T, Nakamura Y. Improvement of Depressed Mood with Green Tea Intake. Nutrients. 2022 Jul 19;14(14):2949.
16. Gilbert, C. The Effect of Tea and Its Constituent L-theanine on Anxiety: A Review of the Literature. (2014).
17. Kimura K, Ozeki M, Juneja LR, Ohira H. L-Theanine reduces psychological and physiological stress responses. Biological psychology. 2007 Jan 1;74(1):39-45.
18. Zhang Y, Tang N, Xia W, Sanjid Seraj S, Pereira M, Velu P, Zhou H, Yang H, Du G. The effect of green tea supplementation on the anthropometric outcomes in overweight and obese women: a time and dose-response meta-analysis of randomized controlled trials. Critical Reviews in Food Science and Nutrition. 2023 May 31:1-0.
19. Wu M, Wu X, Zhu J, Li F, Wei X, Wang Y. Selenium-enriched and ordinary green tea extracts prevent high blood pressure and alter gut microbiota composition of hypertensive rats caused by high-salt diet. Food Science and Human Wellness. 2022 May 1;11(3):738-51.
20. Taati B, Arazi H, Kheirkhah J. Interaction effect of green tea consumption and resistance training on office and

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]

ambulatory cardiovascular parameters in women with high-normal/stage 1 hypertension. The Journal of Clinical Hypertension. 2021 May;23(5):978-86.

21. Nogueira LD, Nogueira Neto JF, Klein MR, Sanjuliani AF. Short-term effects of green tea on blood pressure, endothelial function and metabolic profile in obese prehypertensive women: A crossover randomized clinical trial. Journal of the American College of Nutrition. 2017 Feb 17;36(2):108-15.
22. Li G, Zhang Y, Thabane L, Mbuagbaw L, Liu A, Levine MA, Holbrook A. Effect of green tea supplementation on blood pressure among overweight and obese adults: a systematic review and meta-analysis. Journal of hypertension. 2015 Feb 1;33(2):243-54.
23. Almudhi A, Gabr SA. Green tea consumption and the management of adrenal stress hormones in adolescents who stutter. Biomedical Reports. 2022 Apr 1;16(4):1-9.
24. Wang L, Brennan M, Li S, Zhao H, Lange KW, Brennan C. How does the tea L-theanine buffer stress and anxiety. Food Science and Human Wellness. 2022 May 1;11(3):467-75.
25. Stachowicz M, Lebiedzińska A. The effect of diet components on the level of cortisol. European Food Research and Technology. 2016 Dec;242:2001-9.
26. Silva VL, Mayer MS, Severo AD, Zambra AL, Almeida JM, Sestari LM, Parisi MM, Giacomini AC, Abreu MS, Bortolotto JW, Bonfanti-Azzolin G. L-theanine modulates tissue cortisol levels in zebrafish after single stress. Brazilian Journal of Pharmaceutical Sciences. 2025 Jan 20;61:e23375.
27. Ikram H, Mirza B, Haleem DJ. Neurochemical and behavioral effects of green tea (Camellia sinensis) as observed in animals exposed to restraint stress. Pakistan Journal of Pharmaceutical Sciences. 2017 Mar 1;30(2).
28. Mirza B, Ikram H, Bilgrami S, Haleem DJ, Haleem MA. Neurochemical and behavioral effects of green tea (Camellia sinensis): a model study. Pak. J. Pharm. Sci. 2013 May 1;26(3):511-6.
29. Hodgson AB, Randell RK, Mahabir-Jagessar-T K, Lotito S, Mulder T, Mela DJ, Jeukendrup AE, Jacobs DM. Acute effects of green tea extract intake on exogenous and endogenous metabolites in human plasma. Journal of Agricultural and Food Chemistry. 2014 Feb 5;62(5):1198-208.
30. Higashiyama A, Htay HH, Ozeki M, Juneja LR, Kapoor MP. Effects of l-theanine on attention and reaction time response. Journal of functional foods. 2011 Jul 1;3(3):171-8.
31. Li G, Yang J, Wang X, Zhou C, Zheng X, Lin W. Effects of EGCG on depression-related behavior and serotonin concentration in a rat model of chronic unpredictable mild stress. Food & Function. 2020;11(10):8780-7.
32. Seo DB, Jeong HW, Kim YJ, Kim S, Kim J, Lee JH, Joo K, Choi JK, Shin SS, Lee SJ. Fermented green tea extract exhibits hypolipidaemic effects through the inhibition of pancreatic lipase and promotion of energy expenditure. British Journal of Nutrition. 2017 Jan;117(2):177-86.
33. Elseweidy MM, Abd El-Baky AE, Abdullah A. Effect of green tea catechins intake on brain pattern of certain neurotransmitters in Stz diabetic rats. Res J Med Med Sci. 2009;4:515-8.
34. Zhang L, Xu JY, Du YF, Wang ZM, Li JX, Ou-Yang N, Wang Y, Yin XB, Qin LQ. Green tea and selenium-enriched green tea ameliorates non-alcoholic fatty liver disease through peripheral 5-hydroxytryptamine signals in high-fat diet-fed mice. International Food Research Journal. 2021 Sep 1;28(5):996-1008.

[ Downloaded from nfsr.sbmu.ac.ir on 2026-11-09 ]

[ DOI: 10.66224/nfsr.12.4.33 ]