

# The Association of Food Intake and Physical Activity with Body Composition, Muscle Strength and Muscle Function in Postmenopausal Women

Mohammad Reza Vafa<sup>1\*</sup>, Behnaz Abiri<sup>2</sup>, Sepideh Haghifar<sup>3</sup>, Asma Malkami<sup>3</sup>, Fatemeh Esmaeili<sup>3</sup>, Aida Arefazar<sup>3</sup>, Zohreh Amiri<sup>3</sup>

1- Dept.of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

2- Ph.D. Student in Nutritional Sciences, Dept.of Nutrition, Faculty of Paramedicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

3- National Nutrition and Food Technology Research Institute, Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received: September 2016

Accepted: October 2016

## **A B S T R A C T**

**Background and Objectives:** Sarcopenia is explained as age-related reduction in muscle mass and performance. Some of the most important risk factors for sarcopenia include age, malnutrition, and sedentary life style. The aim of this study was to investigate the association of food intake and physical activity with body composition, muscle strength and muscle function in post-menopausal women.

**Materials and Methods:** In this cross-sectional study, a total of 190 women aged 40-60 years were recruited from the staff working in Iran University of Medical Sciences. Variables consisting of muscle strength and muscle function, anthropometric indices, dietary intakes, and physical activity were evaluated. Statistical analysis was performed using SPSS version 21.

**Results:** There was a positive association between physical activity and handgrip strength (P=0.01), calorie intake and muscle function, protein intake and fat free mass percentage, total fat intake and fat mass percentage (in all associations, P<0.001). In addition, a negative association was observed between carbohydrate intake and muscle function, as well as total fat intake and fat free mass percentage (in both associations, P<0.001). There was not a significant relationship between the other variables (P>0.05).

**Conclusions:** It seems in this middle-aged women group, mild fat and carbohydrate restriction combined with mild or moderate physical activity may lead to reduction in fat mass, increased fat free mass, and improvement in muscle strength and muscle function.

Keywords: Food intake, Physical activity, Body composition, Muscle strength, Muscle function, Postmenopausal women

### Introduction

Advancing age, even in healthy old people, is accompanied by a decrease in muscle mass and muscle strength (1,2). Sarcopenia is a multidimensional age-related disease correlated with age (3), sedentary lifestyle (4), malnutrition (5), and lack of anabolic and anti-catabolic responsiveness to changes in the concentration of extracellular amino acids (6), as well as a rise in abnormal reactive oxygen species (ROS) (7). Hormonal changes in women occur with physiological and behavioral changes (8). Alterations in the size and body composition of postmenopausal women are notable, especially the reduction in lean mass and promotion in fat mass. These alterations also take place in aging, but promote upon the entrance of menopause (8). Recent studies have shown that sarcopenia is an important risk factor for metabolic diseases (9,10). The slow loss of muscle strength leads to functional disability (11,12), the requirement for help in the performance of daily activities (13,14), and elevated

\*Address for correspondence: Mohammad Reza Vafa, Professor of Nutritional Sciences, Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran. E-mail address: rezavafa@yahoo.com

risk of falling and fractures (15). In brief, the functional restrictions and disorders due to sarcopenia decrease the quality of life and functional independence during senescence (4). Therefore, the preservation of the muscle strength in aging is important.

Diet and physical activity are among the steadiest risk factors for sarcopenia and are commonly examined in intervention trials to decrease sarcopenia incidence and/or severity (3). Hence, the present study aimed to investigate the association of food intake and physical activity with body composition, muscle strength and muscle function in middle-aged women.

### Materials and Methods

This cross-sectional study was conducted on middle-aged women staff working in Iran University of Medical Sciences (IUMS), Tehran in 2014. Volunteers could participate in this study if they were menopause female, aged 40-60 years, had a Body Mass Index (BMI) 18.5-29.9, not to be ovariectomized, were not suffering from any chronic or acute diseases, and not taking multivitamins, minerals, laxatives, or hormone medications.

The sample size estimate was based on the correlation between handgrip strength and physical activity. Using the sample size formula for correlation studies and on the basis of the previous studies, the expected minimum correlation coefficient between handgrip strength and physical activity (r=0.20) was estimated (16). We determined that 190 individuals would provide a power of 80% to detect a correlation between these two variables on the basis of two-sided alpha level of 0.05. The 190 women, who met the inclusion criteria, were informed of the aims and protocol of the study, wrote informed consent, and signed participant assent. This study was approved by the Ethics Committee of Iran University of Medical Sciences (IUMS), Tehran, Iran.

Anthropometric, dietary intake and physical activity assessments: We evaluated body weight to the nearest 0.1 kg while the women were minimally clothed and without shoes by e-body (Beurer, Germany). Height was assessed to the nearest 0.5 cm in standing posture, without shoes. Body mass index (BMI) was calculated as body weight (kg) divided by height squared (m<sup>2</sup>). Body composition consisting of Fat Mass percentage (FM%) and Fat Free Mass percentage (FFM%) was taken by e-body (Beurer, Germany).

Dietary intakes were assessed with a 24-h recall for 2 days (1 week day and 1 week-end day), and energy and macronutrient (total fat, carbohydrate, protein) intakes were evaluated using the Nutritionist 4 software. Physical activity level was measured by the Persian and short-form of the International Physical Activity Questionnaire (IPAQ) and demonstrated in Met-Min/week (17).

**Muscle performance:** The upper body strength was assessed by handgrip strength test, and muscle function was measured quantitatively by the use of Time Get Up and Go (TGUG) test.

Handgrip strength was assessed in kg in dominant hand with a hand-held dynamometer (digital hand dynamometer "DIGI-II, Korea"). The women were seated on armchair, while their shoulder adducted and neutrally rotated turned, elbow was flexed to  $90^{\circ}$ , and the forearm and wrist were in a neutral posture (18).

For TGUG test, the women were seated on the chair. They were asked to stand up and walk at common step 3m, then turn to the chair, and sit down again. The obtained value was reported in seconds (s) (19,20).

Statistical analysis was performed by using SPSS version 21.0 (SPSS Inc, Chicago, Illinois) software. The Kolmogorov-Smirnov's test was used to assess the normality of data. Descriptive statistics were expressed as mean $\pm$ SD. Multiple linear regression was used for determining the relationship between food intake and physical activity with body composition, muscle strength and muscle function adjusted for potential confounders. *P* value < 0.05 was considered significant.

#### Results

Descriptive statistics of the participants are shown in Table 1. Dominant hand in 181 (95.3%) women was right hand, and in 9 (4.7%) of them was left hand.

Variable	Value <sup>a</sup>	Min	Max
Age, years	47.3±4.2	43.00	51.50
Height, cm	157.37±6.17	141.00	173.00
Weight, kg	66.09±7.40	50.00	84.70
BMI, kg/m <sup>2</sup>	26.65±2.57	19.50	29.90
Fat mass, (%)	36.94±5.17	26.00	48.50
Fat free mass, (%)	63.04±5.15	51.50	74.00
Energy intake, kcal/day	2200.85±661.44	1082.00	3360.00
Carbohydrate intake, g/day	319.49±110.5	146.70	494.40
Total fat intake, g/day	$71.19 \pm 49.14$	29.44	200.700
Protein intake, g/day	21.24±16.94	22.21	105.40
Physical Activity, Met-min/week	854.76±775.94	33.00	6293.00

Table 1. Characteristics and food intakes of the participants (n=190)

<sup>a</sup> Values are shown as mean±SD.

In linear regression models, after adjustment for physical activity and body weight, there was a positive and significant association between calorie intake and muscle function between protein intake and FFM%, and also between FM% and total fat intake (P<0.001). In addition, a negative and significant association between muscle function and carbohydrate intake, between total fat intake and FFM%, and also between FM% and protein intake was seen (P<0.001). But there was not any significant association between muscle function and other investigated variables of food intake (Table 2).

According to the linear regression analysis and after an adjustment for food intake and body weight, we found a positive and significant association between physical activity and handgrip strength (P=0.01) but not a significant relationship between physical activity and other indicators of body composition and muscle function (Table 3).

Moreover, to evaluate the relationship of body weight with muscle function and muscle strength, after an adjustment for food intake and physical activity, there was a positive and significant association between body weight with FM%, and handgrip strength (P<0.001); however, body weight was negatively associated with FFM% (P<0.001). Furthermore, body weight was not associated with muscle function (P=0.35).

### Discussion

In the present study, we found a positive and significant association between calorie intake and muscle function, protein intake and FFM%, and total fat intake and FM%. Moreover, there was a negative and significant association between muscle function and carbohydrate intake, and between protein intake and FM%. Miller et al. (21) evaluated the relationships between body fat, diet composition, calorie intake and exercise among male (n=107) and female (n=109) adults aged 18-71 years. They found no relationship between energy intake and adiposity, but leanness and exercise were correlated. In fact, they showed that body fat mass is more related to diet composition and physical activity than calorie intake. Valenzuela et al. (22), in a study on 78 men and women aged 60 years, found that low protein intake and insufficient distribution of protein could result to muscle mass reduction and elevated risk of sarcopenia. In the current study, we did not investigate the distribution of protein intake in a day. In addition to physical activity level, calorie and macronutrient intake, several factors such as sex hormone deficiency, vitamin D deficiency, and inflammation are known to be important agents in skeletal muscle structure and function (23) that we did not evaluate them. Furthermore, quality of diet can also affect the body composition and, in turn, muscle performance, which we did not assess them.

[ Downloa
_
_
-
_
4
.nfsr.
9
Ξ.
ā
5
g
0
a,
È.
Š
ž
õ
_
0
_

[ D0I:

	`
8	
극	
ä	
S	
nt	
Da	
<u>:5</u>	
Ŧ	
Da	
g	
Ì	
.=	
nc	
Ξ	
ŭ	
5	
е	
SC	
gu	
<u>_</u>	
nd	
a	
挋	h
u u	
Le	
S	
- B	
SC	
Ĩ	
-	
o	
Ξ	
SO	
a	
8	
0	
ð	,
<u>В</u>	
p	
ar	
Ş	
tal	
Е.	
ğ	
.8	
1 f	
ee	
Ă	
ğ	
<u>ب</u>	
<u>ē</u> .	
at	
Ξ	
SSC	
Ä	
તં	
le	
ab	
Ĥ	

		Fat free r	nass (%)			Fat m	1ass (%)			Handgrip str	ength (kg)			Muscle fu	inction (s) <sup>a</sup>	
	Mode	11*	Mode	il 2 **	W	odel 1	Mode	31 2	Mc	del 1	Model	2	Mo	del 1	Model	2
I	β	Ь	β	Ь	β	Ь	β	Ь	β	Ч	β	Ч	β	Ь	β	Ь
Calorie intake, kcal/day	-0.51	<0.001	-0.45	<0.001	0.06	0.51	0.45	0.06	-0.34	<0.001	0.05	0.77	0.31	<0.001	1.16	<0.001
CI (95%)	(-0.005-	-0.003)	(-0.00	(00.0-2)	(0.003	-0.005)	(0.00-	0.007)	(-0.006-	-0.002)	(-0.004-	0.005)	(0.001-	-0.002)	(0.002-0	(800)
Carbohydrate intake, g/day	-0.26	<0.001	-0.11	0.45	0.26	<0.001	0.10	0.47	-0.11	0.13	-0.08	0.46	0.08	0.22	-3.33	<0.001
CI (95%)	(-0.019-	-0.006)	(-0.01	(600.0-6	(0.006	-0.019)	(-0.09	-0.019)	(-0.018	- 0.002)	(-0.21-	0.1)	(-0.001	-0.006)	(-0.025	0.006)
Total fat intake, g/day	-0.50	<0.001	-3.57	<0.001	0.50	<0.001	3.57	<0.001	-0.45	<0.001	-0.52	0.56	0.27	<0.001	-2.60	0.08
CI (95%)	-0.067-	-0.041)	(-0.617	0.133)	(0.041	-0.067)	(0.132-	-0.616)	(-0.093-	-0.052)	(-0.364-(	(199)	(0.008-	0.024)	(-0.322-	0.020)
Protein intake, g/day	-0.22	<0.001	0.27	<0.001	0.22	<0.001	-0.27	<0.001	-0.05	0.42	0.05	0.43	0.14	0.04	0.05	0.65
CI (95%)	(-0.112-	-0.027)	(0.027	7-0.141)	(0.026	-0.112)	(-0.141	0.26)	(-0.093	- 0.039)	(-0.04-0	.093)	(0.001-	-0.049)	(-0.031-	0.050)
<sup>*</sup> Unadjusted **Adjusted for physic <sup>a</sup> Muscle function wa	al activity and s measured by	1 weight / use of Time	Get Up and	Go (TGUG) te	st											

õ	
:19	
_n=	
ts (	
an	
cip.	
urti.	
ğ	
the	
ш.	
U	
cti	
nn	
еf	
scl	
nu	
ı pı	
ar	
ŝth	
eng	
str	
cle	
nsc	
m	
л,	
iti	
SOC	
li	
00	
dy	
pq	
nd	
y a	
Ϋ́τ.	
Ę	
la	
ica	
ıys	
þ	
Sen	
M	
bet	
uo	
ati	
) C	
SS	
A.	
le3	
abl	
Ë	

		Fat free	: mass (%)			Fat m	18S (%)			Handgrip s	trength (k	(g)		Muscle fu	nction (s)	
		Model 1 <sup>*</sup>	Mode	12**	Mc	odel 1	Mod	el 2	M	odel 1	Mode	el 2	Mc	odel 1	Mod	el 2
	В	Р	β	Ь	β	Ь	β	Ь	β	Ь	β	Ь	β	Ь	β	Ь
Physical activity, (Met-min/week)	0.38	<0.001	-0.19	0.39	-0.38	<0.001	0.20	0.38	0.41	<0.001	0.43	0.01	-0.22	<0.001	0.55	0.057
CI (95%)	(0.0)	02-0.003)	(-0.004-	0.002)	(-0.003-	-0.002)	(-0.002-(	0.004)	(0.003	-0.005)	(0.001-	0.008)	(-0.001)	- 0.00)	-00.0)	.004)
*Unadjusted **Adjusted for food intak	e and bod	y weight														

Our findings demonstrated no significant association between physical activity level, and FM% and FFM%. Westerterp et al. (24) investigated the relationship between physical activity level and parameters of aging among the 136 women and 180 men over 20 years. They found that advancing age was correlated with lower activity levels and lower fat free mass. After controlling for age, they found no association between physical activity level and fat free mass. In old subjects with relatively high physical activity level compared with those with low levels of physical activity, there was no difference in fat mass and fat free mass (24). This study found that the effect of exercise on body composition depends on the kind of exercise (24). Resistance exercise elevates fat free mass by 1.1 to 2.1 kg, while aerobic exercise had no impact on fat free mass. In addition, in older adults, exercise training did not lead to a promotion in TEE (Total Energy Expenditure), and exercise activity was compensated by reduction in spontaneous activity (24).

In the current study, there was a significant relationship between physical activity level and handgrip strength. Kuh *et al.* (25) revealed that middle-aged women had much poorer handgrip strength, weaker balance time, and poorer chair rise time, were less active and suffered from health problems. We found no significant association between physical activity level and muscle function, which is probably resulted from lack of association between physical activity level, and FM% and FFM%.

In spite of these findings, it should be noted that, because of the cross-sectional nature of this study, causality could not be determined, and could not find the temporality of the observed results. In this regard, life-span (longitudinal and prospective) studies are needed.

There were some limitations in the present study; including the cross-sectional design of this study, and lack of assessment of oxidative stress and inflammatory biomarkers in the participants. To evaluate the muscle strength, we measured the upper body muscle strength, only.

In conclusion, it seems in this middle aged women group, mild fat and carbohydrate restriction combined with mild or moderate physical activity may lead to reduction in fat mass, and increased fat free mass and also improvement in muscle performance.

### Acknowledgement

We thank the staff of Iran University of Medical Sciences for their cooperation and participation in this study.

### **Financial disclosure**

None conflict of interest has been declared.

### Financial disclosure

The authors declared no financial interest.

### **Funding/Support**

This work was financially supported by the Vice Chancellor of Research, Iran University of Medical Sciences (Tehran, Iran).

### References

- 1. Lexell J. Human ageing, muscle mass, and fiber type composition. J Gerontol 1995;50A:11–6.
- 2. Hurley BF. Age, gender, and muscular strength. J Gerontol 1995; 50A:41–4.
- Beasley JM, Shikany JM, and Thomson CA. The role of dietary protein intake in the prevention of sarcopenia of aging. Nutr Clin Pract. 2013 December; 28(6): 684–690.
- 4. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. J Am Geriatr Soc 2002;50:889–96.
- 5. Bos C, Benamouzig R, Bruhat A, Roux C, Mahe S, Valensi P, et al. Short-term protein and energy supplementation activates nitrogen kinetics and accretion in poorly nourished elderly subjects. Am J Clin Nutr 2000;71:1129–37.
- 6. Katsanos CS, Kobayashi H, Sheffield-Moore M, Aarsland A, Wolfe RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. Am J Physiol Endocrinol Metab 2006;291:E381–7.
- 7. Fulle S, Protasi F, Tano GD, Pietrangelo T, Beltramin A, Boncompagni S, et al. The contribution of reactive oxygen species to sarcopenia and muscle ageing. Experimental Gerontology. 2004: 39.1: 17-24.
- L. Dennerstein, Depression in the menopause, Obstet. Gynaecol. Clin. N. Am. 1987: 4: 33–39.
- Metter EJ, Talbot LA, Schrager M, Conwit R. Skeletal muscle strength as a predictor of all-cause mortality in healthy men. J Gerontol A Biol Sci Med Sci 2002;57:B359e65.
- 10. Kim MK, Baek KH, Song KH, Il Kang M, Park CY, Lee WY, et al. Vitamin D deficiency is associated with sarcopenia in older Koreans, regardless of obesity: the Fourth Korea National Health and Nutrition Examination Surveys (KNHANES IV) 2009. J Clin Endocrinol Metab 2011;96:3250e6.

- 11. Samson MM, Meeuwsen IBAE, Crowe A, Duursma SA, Verhaar HJJ. Relationships between physical performance measures, age, height and body weight in healthy adults. Age Ageing 2000;29: 235–42.
- 12. Bassey EJ, Fiatarone MA, O'Neill EF, Kelly M, Evans WJ, Lipsitz LA. Leg extension power and functional performance in very old men and women. Clin Sci 1992;82:321–7.
- Avlund K, Schroll M, Davidsen M, Løvborg B, Rantanen T. Maximal isometric muscle strength and functional ability in daily activities among 75-year-old men and women. Scand J Med Sci Sports 1994;4:32–40. 15.
- Hyatt RH, Whitelaw MN, Bhat A, Scott S, Maxwell JD. Association of muscle strength with functional status of elderly people. Age Ageing 1990;19:330–6.
- 15. Wolfson L, Judge J, Whipple R, King M. Strength is a major factor in balance, gait, and the occurrence of falls. J Gerontol 1995;50:64–7.
- 16. Katzmarzyk P.T, Craig C.L, & Gauvin L. Adiposity, physical fitness and incident diabetes: the physical activity longitudinal study. Diabetologia 2007; 50:538–544.
- Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. Med Sci Sports Exerc 2003; 35:1381–95.
- Budziareck MB, Pureza Duarte RR, Barbosa-Silva MC. Reference values and determinants for handgrip strength in healthy subjects. Clin Nutr 2008; 27:357–62.
- 19. Samson MM, Meeuwsen IB, Crowe A, Dessens JA, Duursma SA, Verhaar HJ. Relationships between

physical performance measures, age, height and body weight in healthy adults. Age Ageing 2000; 29:235–42.

- 20. Wall JC, Bell C, Campbell S, Davis J. The Timed Getupand-Go test revisited: measurement of the component tasks. J Rehabil Res Dev 2000; 37:109–13.
- Miller WC, Lindeman AK, Wallace J, and Niederpruem M. Diet composition, energy intake, and exercise in relation to body fat in men and women. Am J Clin Nutr 1990: 52: 426-30.
- 22. Valenzuela RE, Ponce JA, Morales-Figueroa GG, Muro KA, Carreón VR, Alemán-Mateo H. Insufficient amounts and inadequate distribution of dietary protein intake in apparently healthy older adults in a developing country: implications for dietary strategies to prevent sarcopenia. Clinical Interventions in Aging 2013:8: 1143–1148.
- 23. Park S, Ham J, Lee BK. A positive association of vitamin D deficiency and sarcopenia in 50 year old women, but not men. Clinical Nutrition 33 (2014) 900e905.
- 24. Westerterp KR and Meijer EP. Physical Activity and Parameters of Aging: A Physiological Perspective. Journals of Gerontology: SERIES A 2001: 56A (Special Issue II):7–12.
- 25. Kuh D, Bassey EJ, Butterworth S, Hardy R, Wadsworth ME, and the Musculoskeletal Study Team. Grip strength, postural control, and functional leg power in a representative cohort of British men and women: Associations with physical activity, health status, and socioeconomic conditions. J Gerontol A Biol Sci Med Sci 2005: 60A: 2,224–231.