



Persian Gum: A Novel Natural Hydrocolloid

Soleiman Abbasi *

Food Colloids and Rheology Lab., Department of Food Science and Technology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

Received: November 2016

Accepted: December 2016

ver the past few decades, scientists have paid special attention to studying novel and natural hydrocolloids due to the growing demand for ready meals, public awareness on the importance of natural fibers in daily diets, and their wide range of application, especially in foods, pharmaceuticals and herbal medicine. Moreover, natural hydrocolloids are preferred to biotechnologic ones (e.g., gellan, xanthan and curdlan) as they are non-toxic. natural. biodegradable, usually biocompatible and ecologically friendly. Persian gum (PG) is one of these natural hydrocolloids, which has been introduced and studied on mostly by the author and some other researchers during the past few years. The author attempted to name it Persian gum, representing its origin the Persia or Iran, as the competitor of the very well-known "gum Arabic", the exudate of acacia trees (Acacia senegal or Acacia Its botanical source, physicochemical, seval). structural. rheological, functional properties, interaction with other macromolecules (proteins and polysaccharides), and possible applications in foodstuffs have been extensively reviewed (1-4).

Amygdalus scoparia Spach (Synonym: Prunus scoparia Spach) is one of the wild (mountain) almond species, which can be found along with other wild species mostly in semi-arid areas of the central Asia, Irano–Tourani and Zagrosi regions. It is appreciated mostly due to its role in the economy (the usage of its by-products in perfume, pharmaceutical and chemical industries), as a natural resource, as well as in soil erosion control. For those reasons, it is nowadays being widely used in breeding programs (to generate rootstocks that are adaptable to climate changes) and planting (seeds and seedlings) in mountains and deserts to control the rapidly happening desertification process in Iran (1).

These features make it very clear that *Amygdalus scoparia* Spach can be considered as a reliable and sustainable source for providing a natural exudate gum, PG, along other products (bitter almond, kernel oil, timber, and ornamentals). Iran, owing to its geographical and environmental situation, has traditionally been a great source for this product such that, at present, it annually exports over 400,000 kg of PG. It is predicted that this number will tremendously increase in a matter of a few years or so once the artificial population of seedlings is established (1, 3).

The growing demand in food and pharmaceutical industries for safe, low price, easily accessible and biodegradable, sustainable, biocompatible, ecologically friendly and renewable natural sources for stabilizers and emulsifiers has persuaded scientists to investigate the potential capabilities of existing natural gums and mucilages, as alternatives for the current ones, which are getting more and more expensive (e.g., gum tragacanth), unsustainable (e.g. gum Arabic) due to the periodical crisis in the producing countries like Senegal, Sudan and Somalia, or with health concerns (e.g., bio-based gums). To this end, PG is a relatively unknown natural hydrocolloid but with abundant production and countrywide distribution, so that it has been considered as a potential alternative, and its applicability characteristics and have been extensively studied (1).

PG, in its natural state, is used as herbal remedy (*e.g.*, as a poultice for swollen joints), an antiparasite, teeth pain healer, appetizer, anti-cough agent, hair conditioner, and skin glazer in Iran and

some other countries (1–4). Apart from these traditional applications, it has other potentials in stabilization of acidic milk-based drinks (5–8), flixweed seeds drink (9), orange peel essential oil nanoemulsions (10), partial replacer of gelatin in formulation of jelly or gummy candies (11), edible films (12), dairy products, emulsions, tomato ketchup, mayonnaise, salad dressings, milk and dark chocolate, bakeries, low-fat cheese, and many other applications (*e.g.*, in personal communications).

All in all, it is a competitively low cost, novel, and natural hydrocolloid, which can potentially be used as a desirable substitute for microbial or other natural but expensive plant hydrocolloids. However, its partial solubility (70 %w/w insoluble) is a major challenge, particularly when it is needed to be used in liquid or semi-liquid formulations (foods, pharmaceuticals, and cosmetics) or as an emulsifier for which fractionation (separation of the soluble and insoluble parts) is a must. Therefore, further investigation of its chemical characterization as well as the solubilisation of its insoluble fraction by various methods is needed (13-14). This could improve and expand its functionality. Furthermore, the preservative function of special fractions of PG, as a natural hydrocolloid, also needs to be investigated.

Financial disclosure

The author declare no financial interest.

References

- 1. Abbasi S. Challenges towards characterization and application of a novel hydrocolloid. Current Opinion in Colloid and Interface Science 2017; Accepted.
- 2. Dabestani M, Kadkhodaee R, Phillips GO, Abbasi S. Persian gum: A comprehensive review on its physicochemical and functional properties. Food Hydrocolloids, 2017; Revised.
- 3. Abbasi S, Rahimi S. Persian gum. In: Mishra S, editor. Encyclopedia of Biomedical Polymers and Polymeric

Biomaterials. USA: Taylor & Francis Group LLC; 2015, pp. 9001–9011.

- Fadavi G, Ghiasi M, Zargarran A, Mohammadifar M.A. Some physicochemical and rheological properties of Zedo (Farsi) gum exudates from Amygdalus scoparia. Nutrition and Food Sciences Research 2017; 4: 31–38.
- 5. Abbasi S. Mohammadi S. Stabilization of milk–orange juice mixture using Persian gum: Efficiency and mechanism. Food Bioscience 2013; 2: 53–60.
- 6. Teimouri S, Abbasi S, Scanlon MG. Stabilization mechanism of various inulins and hydrocolloids: Milk– sour cherry juice mixture. International Journal of Dairy Technology 2017; doi: 10.1111/1471-0307.12376.
- 7. Azarikia F, Abbasi S. Mechanism of soluble complex formation of milk proteins with native gums (tragacanth and Persian gum). Food Hydrocolloids 2016; 59: 35–44.
- Teimouri S, Abbasi S, Sheikh N. Effects of gamma irradiation on some physicochemical and rheological properties of Persian gum and gum tragacanth. Food Hydrocolloids 2016; 59: 9–16.
- Behbahani MS, Abbasi S. Stabilization of Flixweed (Descurainia sophia L.) syrup using native hydrocolloids. Iranian Journal of Nutrition Sciences & Food Technology 2014; 9: 31–38. [In Persian]
- Mirmajidi Hashtjin A, Abbasi S. Nano-emulsification of orange peel essential oil using sonication and native gums. Food Hydrocolloids 2011; 44: 40–48.
- 11. Abbasi S, Mohammadi S, Rahimi S. Partial substitution of gelatin with Persian gum and use of olibanum in production of functional pastille. Iranian Journal of Biosystems Engineering 2011; 42: 121–131. [In Persian]
- 12. Khalighi S, Abbasi S. Film forming properties of Persian gum. Iranian Journal of Food Science and Technology, Accepted. 2017. [In Persian]
- Mohammadi S, Abbasi S, and Scanlon MG. Development of emulsifying property in Persian gum using octenylsuccinic anhydride (OSA). International Journal of Biological Macromolecules 2011; 89: 396– 405.
- 14. Samari Khalaj M, Abbasi S. Influence of chemical modification on solubility of insoluble fraction of Persian gum. Journal of Research and Innovation in Food Science and Technology 2014; 3: 171–184. [In Persian]