



Original Article

The Effects of Chicory Leaf Aqueous Extract on Body Weight, Serum Glucose and Lipid Levels in Streptozotocin Induced Diabetic Rats

Reza Gorjipour¹, Jan-Mohamad Malekzadeh^{2*}, Haibatollah Sadeghi², Jamshid Mohammadi², Fariba Malekzadeh³

1- Student Research Committee, Yasuj University of Medical Sciences, Yasuj, Iran.

2- Medicinal Plants Research Center, Yasuj University of Medical Sciences, Yasuj, Iran.

3- M.Sc. in English Language Teaching, Boyer Ahmad Department of Education, Yasuj, Iran.

Received: June 2017

Accepted: August 2017

ABSTRACT

Background and Objectives: Diabetes mellitus is a common chronic disease worldwide. Although it is treated with a number of methods including diet therapy, hypoglycemic agents, insulin and some herbs, its' prevalence and complications are rapidly increasing. *Chicory intybus* has been used as a traditional diabetes treatment in Iran, Egypt, and other countries but there is a lack of convincing evidence on its effect. This study was designed to evaluate the effects of aqueous extract of chicory leaf on body weight, serum insulin, glucose and lipids in diabetic male Wistar rats.

Materials and Methods: Forty male Wistar rats were divided into five groups. Four groups were given streptozotocin intraperitoneally to induce diabetes. Three of the diabetic groups received varying concentrations of chicory aqueous extract for three weeks (12, 25 and 50 mg /kg body weight). The forth diabetic group and the non-diabetic group received distilled water (diabetic and healthy control groups respectively). Fasting blood glucose, serum triglyceride (TG), total cholesterol (TC), LDL and HDL levels were measured after 3 weeks of treatment.

Results: The results did not show any significant effects on fasting blood glucose (173.5 ± 126 , 154.5 ± 100 , 247.7 ± 46 , 170.7 ± 150.8 mg/dL) and serum lipids; LDL (19.5 ± 7.7 , 26.4 ± 6.7 , 24.2 ± 7.7 , 24.7 ± 6.8 mg/dL), HDL (41 ± 22.8 , 41.2 ± 12 , 44.7 ± 12.1 , 40 ± 4.6 mg/dL), TG (103.5 ± 35 , 80.4 ± 53.8 , 111 ± 70 , 94 ± 8.9 mg/dL) and TC (76.2 ± 27.3 , 80.4 ± 53.8 , 77 ± 18.4 , 72.7 ± 15 mg/dl) in Streptozotocin induced diabetic groups receiving 12, 25 and 50 mg/kg aqueous chicory extract and diabetic control groups respectively.

Conclusions: Chicory leaf aqueous extract had no significant effects on serum glucose and lipids of streptozotocin-induced diabetic rats.

Keywords: Diabetes, Chicory extracts, Blood glucose, Cholesterol, Triglyceride, Rat

Introduction

Diabetes mellitus is a very common health problem of the world, in both developed and developing countries and is one of the leading causes of death worldwide (1). It is estimated that the number of world diabetics will exceed to over 400 million by 2030 (2). Recent studies indicate that the prevalence of diabetes in Iran, and in other developed and developing countries is increasing (3-6). Diabetes results in severe complications including hyperlipidemia, neuropathy, nephropathy, retinopathy

and cardiovascular disorders and therefore impose great economic costs to the families and countries (7, 8). Diet therapy, exercise, pharmacotherapy and herbal medicines are strategies to treat diabetes mellitus. Herbs have been used for diabetes mellitus treatment in ancient Iran, China, Egypt and other countries (9, 10). Chicory intybus (*Cichorium intybus* L.), is a traditional herbal medicine, which belongs to the *Asteraceae* family with six species mostly in Europe and Asia (11) also is known as *Kasni* in

Iran(12). This herb is said to have a number of beneficial and therapeutic properties such as its use in the treatment of wound healing(13), depression(14), hypertension(15) jaundice (16), hiccups (17) and diabetes (18). The effects of herbs on diabetes treatment is currently the subject of vast number of studies (9, 19-26). Alkaloids, inulin, sesquiterpene lactones, coumarins, unsaturated sterols, flavonoids, saponins, tannins and other medicinal constituents, potentially effective in diabetes treatment, are detected in the chicory roots and leaves (27). The effects of ethanol extract of whole plant *Cichorium intybus* on diabetes was studied in STZ diabetic rats and were reported to reduce blood glucose, blood triglyceride and blood cholesterol (28). Ziamajidi *et al.* also reported hepatic protective effects for chicory extract in hepatic steatosis in rats (*in vivo*), and in HepG2 cells (*in vitro*) (29). The components of chicory root are different from leaves and so their effects on diabetes treatment may be different. The effects of chicory whole plant, chicory seed (29) and chicory root on blood glucose and lipids have been reported previously (28-30). However, the effects of aqueous extract of chicory leaf on weight, blood glucose and lipids need to be examined. This study was performed to evaluate the effects of chicory leaf aqueous extract on blood glucose and lipids in Streptozotocin (STZ) induced diabetic rats.

Materials and Methods

Forty adult male Wistar rats, 12 weeks of age, have randomly divided into five groups of eight. Four groups of rats were induced diabetes by intraperitoneal injection of 50 mg/kg of STZ. The fifth group of rats did not receive STZ (healthy control). A diabetic group (diabetic control) received distilled water through gavage during the study. The third, fourth and fifth groups of diabetic rats received chicory leaf extract through the gavage (12.5, 25 and 50 milligrams/kilogram body weight, respectively).

The rats were kept in $20 \pm 2^{\circ}\text{C}$ room temperature, 12 h light/dark cycle and had free access to standard rat diet and water.

Diabetes induction: Diabetes was induced by intraperitoneal injection of 50 mg/kg body weight STZ. Blood glucose was measured using a glucometer (high precision German IME-DC) at intervals of 24 hours during the first week, and if blood glucose was

above 200 mg/dl, rats were considered as diabetic (31).

Extraction: The *Chicory intybus* was collected from the margins of dry lands farming in Boirahmad, Iran. Plant leaves were detached from roots, washed gently, dried under the shadow, and powdered using a kitchen mill. One kilogram of powdered leaves was soaked in 5 liters of distilled water and was stored at room temperature for 72 hours. The soaked mixture was stirred and filtered using Whatman filter paper No 1. The dissociated liquid filtrate was concentrated using rotary apparatus under the conditions of vacuumed atmosphere and dried at 50°C temperature. The solutions were made using dried extract according to the required concentrations.

Blood sampling and laboratory tests: After diabetes induction, chicory extract solutions were given to the rats through daily gavage for three weeks. Blood glucose of all rats were measured at the baseline (day 1), two weeks (day 14) after receiving chicory extract and at the end of the study on the last day of third week (day 21), using a high precision German IME-DC glucometer in rat tail blood.

Finally, after an overnight fasting, rats were anesthetized with *ether du petrol* and one milliliter blood was drawn from each rat's heart. Blood samples were centrifuged by 3000 RPM, the sera were separated and sent to the laboratory to measure blood glucose, serum cholesterol, LDL-C, HDL-C and triglyceride. The weight of the rats were measured at the 1st and last day of the intervention.

Data Analysis: Findings were analyzed using the SPSS version 19. Statistical significance was detected at *p* value less than 0.05. Differences between groups were tested using one way analysis of variance followed by *Scheffe post hoc* test to determine significant difference. Repeated measure ANOVA were used to test the fasting blood glucose changes in three consecutive measures. Before-after groups' mean weight were compared with paired samples *t* test.

Results

The mean weight of all groups at the baseline measurements were not significantly different. The animals final body weights (day 21) showed a significant reduction from the initial weights in all diabetic groups but the healthy control group did not show any significant weight change. The final body

weight between the diabetic groups were not significantly different (Table 1).

Increasing fasting serum glucose in diabetic groups receiving the STZ compared with the healthy control group showed that the rats were indeed diabetic ($F=7.74$; $p<0.001$). Serum fasting glucose levels in diabetic groups were not significantly different after two weeks ($F=0.08$; $p=0.96$), and after

three weeks of extract gavage ($F=0.48$; $p=0.7$), as shown in table 2. Table 3 shows serum lipid levels of all groups. As shown, there is a clear, although non-significant difference between the lipid levels of healthy control group with diabetic groups. Serum levels of triglyceride are increased in diabetic groups but cholesterol, HDL-C, and LDL-C levels are decreased compared to the healthy control.

Table 1. Effects of chicory extract on body weight levels (gr).

Group	Time	Initial	Final	Weight change	Paired t test sig
Healthy control		409.1±48	410±48.6 ^ψ	0.83±7.3 ^ψ	0.79
Diabetic control		335±75.9	247±36.8*	-88±40*	0.022
Diabetic +12.5 mg/kg extract		365.7±45.1	294.3±26.4*	-71.4±30.8*	0.001
Diabetic +25 mg/kg extract		346.7±58.6	265±25.3*	-81.7±42.5*	0.001
Diabetic +50 mg/kg extract		390±40.8	288.7±14.4*	-101.2±27.8*	0.005
ANOVA,		F=1.6	F=18.5	F=9.7	
Sig		p=0.2	P=0.0001	P=0.0001	

The values are mean±SD for 8 rats in each group. Means with different superscript (*, ψ) within a column are significantly different from each other at $P<0.05$ as determined by one way ANOVA and Scheffe post hoc test.

Table 2. Mean and standard deviation of blood glucose (mg/dl) in all groups on the baseline, two weeks and three weeks of study

Group	Time	Baseline	Day 14	Day 21	P Value
Healthy control		62.6±9.3 ^ψ			
Diabetic control		308.7±133.8*	168.7±91	173.5±126	F=2.8, P=0.09 ANOVA Repeated measure
Diabetic +12.5 mg/kg extract		202±71*	175.8±87	154.5±100	
Diabetic +25 mg/kg extract		259.6±76*	204.3±129	247.7±46	
Diabetic +50 mg/kg extract		190.2±150*	176.2±104.7	170.7±150.8	
F ANOVA		F=7.74	F=0.08	F=0.48	
P Value		p<0.001	P=0.96	p=0.7	

The values are mean±SD for 8 rats in each group. Means with different superscript (*, ψ) within a column are significantly different from each other at $P<0.05$ as determined by Scheffe post hoc test

Table 3. Mean and standard deviation of serum lipids levels (mg/dl) in groups of rats at the end of the experiment

Group	Factors	TC	HDL	LDL	TG
Healthy control		90.8±18.4	57.7±12.6	30±6.7	53±10.7
Diabetic control		76.2±27.3	41±22.8	19.5±7.7	103.5±35
Diabetic +12.5 mg/kg extract		80.4±53.8	41.2±12	26.4±6.7	80.4±53.8
Diabetic +25 mg/kg extract		77±18.4	44.7±12.1	24.2±7.7	111±70
Diabetic +50 mg/kg extract		72.7±15	40±4.6	24.7±6.8	94±8.9
P Value		F=0.59 P=0.63	F=1.3 p=0.3	F=1.47 p=0.25	F=1.06 p=0.4

The values are mean±SD for 8 rats in each group.

Discussion

In this study, we evaluated the effects of chicory leaf aqueous extract on blood glucose and lipids level of STZ induced diabetic rats. Our findings did not show any significant effects of chicory leaf aqueous extract on serum glucose and lipids in STZ induced diabetic rats. The outcome of the current study is contrary to Pushparaj *et al.*, (28) that in their study on the effects of alcoholic extract of chicory, a 20 percent reduction in serum glucose, 91 percent in triglycerides and 16 percent reduction in the serum cholesterol of STZ-induced diabetic rats was reported. Pushparaj and colleagues used whole plant including leaves, roots and stems for alcoholic extraction and also a higher concentrations of extraction than ours (120 mg/kg vs. 12.5, 25 and 50 mg/kg in our study) that may explain the different findings in two studies. The root and stems has different compounds than leaves, especially soluble fibers. Inulin a major component of chicory root is a fructan and soluble fiber and has hypolipidemic effects through the growth of bifidobacteria and reducing fat absorption(32).

The STZ diabetes induced rats showed a significant weight reduction compared to the control group, but Chicory leaf extract did not affect the weight of the rats or did not prevent weight reductions. Ghamarian *et al* showed that chicory seed extract prevents weight reduction in both STZ diabetic rats and Niacinamid-STZ diabetic rats (33). Ghamarian *et al* used the seed extracts injection intraperitoneal, but we used leave extract by gavage. The differences in plant part (leave against seeds) and application route (gavage against intraperitoneal injection) may explain the inconsistencies shown between the two studies.

Kim *et al* (32) had examined antilipidemic properties of aqueous extract of chicory root and showed that healthy rats receiving chicory extract had higher Serum HDL cholesterol concentrations, but reported no significant effect on cholesterol concentrations.

Khaksari and colleagues studied the effects of chicory containing diet on the blood glucose in diabetic rats. It was shown that blood glucose was decreased significantly in rats in a dose-dependent manner(30). Touseh and colleagues examined the effect of chicoric acid, a component in chicory plant, on blood glucose of diabetic rats. Their findings

showed that chicoric acid has the potential anti-diabetic and insulin secretagogue effect (34).

Previous studies used whole chicory plant including the leaf, root and stem (28) or the root extracts (32) or used the whole plant mixed with the rats diet(30), but we used the aqueous extract of chicory leaves. Our findings on chicory leaf extract is not similar to other studies, which may be due to the different application route and compounds in different parts of chicory. Chicory has abundant amounts of inulin in the root(35) and contain anthocyanins, phenols (36) as well as sesquiterpenes (37). Inulin is a water-soluble fiber and possesses hypolipidemic effects (38). Based on our findings chicory leaf aqueous extract is not effective in reducing blood glucose or blood lipids.

Acknowledgement

The authors would like to thank the Medicinal Plants Research Center/deputy of Research and technology of Yasuj University of Medical Sciences for funding this research. We also thanks Mr. Reza Mohamadi, the laboratory technician in herbal medicine research center for his assistance in extract preparation.

Financial disclosure

All experimental procedures including treating the rats used in this study were conducted based on Ethical regulation of medical research on laboratory animals in Iran. The study was approved by ethic committee of Yasuj University of Medical Sciences.

Funding/Support

This research was approved and funded by deputy of Research and technology of Yasuj University of Medical Sciences with the grant number 11316.

References

1. Nanditha A, Ma RC, Ramachandran A, Snehalatha C, Chan JC, Chia KS, et al. Diabetes in Asia and the Pacific: Implications for the Global Epidemic. *Diabetes Care*. 2016;39(3):472-85.
2. Chen L, Magliano DJ, Zimmet PZ. The worldwide epidemiology of type 2 diabetes mellitus--present and future perspectives. *Nature reviews Endocrinology*. 2012;8(4):228-36.
3. Sadeghi M, Roohafza H, Shirani S, Poormoghadas M, Kelishadi R, Baghaei A, et al. Diabetes and associated cardiovascular risk factors in Iran: the Isfahan Healthy

- Heart Programme. Annals of the Academy of Medicine, Singapore. 2007;36(3):175-80.
4. Larejani BZ, F. Epidemiology of diabetes melitus in Iran. Iranian Journal of Diabetes and Metabolism. 2001;1(1):1-8.
5. Larijani B, Abolhasani F, Mohajeri-Tehrani M, Tabatabaie O. Prevalence of diabetes mellitus in Iran in 2000. Iranian Journal of Diabetes and Metabolism. 2005;4(3):75-83.
6. Safari M, Yazdanpanah B, Yazdanpanah B, Mobasheri A. A population-based screening of type 2 diabetes in high-risk population of Yasuj, Iran. Journal of health, population, and nutrition. 2014;32(4):677-86.
7. Nentwich MM, Ulbig MW. Diabetic retinopathy - ocular complications of diabetes mellitus. World journal of diabetes. 2015;6(3):489-99.
8. Viswanathan V. Preventing microvascular complications in type 1 diabetes mellitus. Indian journal of endocrinology and metabolism. 2015;19(Suppl 1):S36-8.
9. Ko CH, Yi S, Ozaki R, Cochrane H, Chung H, Lau W, et al. Healing effect of a two-herb recipe (NF3) on foot ulcers in Chinese patients with diabetes: a randomized double-blind placebo-controlled study. Journal of diabetes. 2014;6(4):323-34.
10. Kibiti CM, Afolayan AJ. Herbal therapy: A review of emerging pharmacological tools in the management of diabetes mellitus in Africa. Pharmacognosy Magazine. 2015;11(Suppl 2):S258-S74.
11. Street RA, Sidana J, Prinsloo G. Cichorium intybus: Traditional Uses, Phytochemistry, Pharmacology, and Toxicology. Evidence-based complementary and alternative medicine : eCAM. 2013;2013:579319.
12. Nasiri A, Ziamajidi N, Behrouj H, Abbasalipourkabir R, Dehghan A. The effects of aqueous extract of chicory root on steatosis, lipid profile and liver damage enzyme markers in tamoxifen-treated rats. Molecular and Biochemical Diagnosis (Journal). 2014;1(3):185-94.
13. Suntar I, Kupeli Akkol E, Keles H, Yesilada E, Sarker SD, Baykal T. Comparative evaluation of traditional prescriptions from Cichorium intybus L. for wound healing: stepwise isolation of an active component by in vivo bioassay and its mode of activity. Journal of ethnopharmacology. 2012;143(1):299-309.
14. Tavakkoli-Kakhki M, Eslami S, Motavasselian M. Nutrient-rich versus nutrient-poor foods for depressed patients based on Iranian Traditional Medicine resources. Avicenna journal of phytomedicine. 2015;5(4):298-308.
15. Ahmad L, Semotiuk A, Zafar M, Ahmad M, Sultana S, Liu QR, et al. Ethnopharmacological documentation of medicinal plants used for hypertension among the local communities of DIR Lower, Pakistan. Journal of ethnopharmacology. 2015;175:138-46.
16. Amiri MS, Joharchi MR, Taghavizadehyazdi ME. Ethno-medicinal plants used to cure jaundice by traditional healers of mashhad, iran. Iranian journal of pharmaceutical research : IJPR. 2014;13(1):157-62.
17. Mohammadi Q, Minae MB, Somi MH, Mosaddegh M, Kamalinejad M. Novel use of chicory for the treatment of hiccups in liver obstruction: in Iranian traditional medicine. Iranian Red Crescent medical journal. 2013;15(11):e6647.
18. Judžentienė A, Būdienė J. Volatile constituents from aerial parts and roots of Cichorium intybus L. (chicory) grown in Lithuania. Chemija. 2008;19:25-8.
19. Akash MS, Rehman K, Chen S. Spice plant Allium cepa: dietary supplement for treatment of type 2 diabetes mellitus. Nutrition. 2014;30(10):1128-37.
20. Boaduo NK, Katerere D, Eloff JN, Naidoo V. Evaluation of six plant species used traditionally in the treatment and control of diabetes mellitus in South Africa using in vitro methods. Pharmaceutical biology. 2014;52(6):756-61.
21. Ghosh S, More P, Derle A, Patil AB, Markad P, Asok A, et al. Diosgenin from Dioscorea bulbifera: novel hit for treatment of type II diabetes mellitus with inhibitory activity against alpha-amylase and alpha-glucosidase. PloS one. 2014;9(9):e106039.
22. Li Y, Li JJ, Wen XD, Pan R, He YS, Yang J. Metabonomic analysis of the therapeutic effect of Potentilla discolor in the treatment of type 2 diabetes mellitus. Molecular bioSystems. 2014;10(11):2898-906.
23. Medagama AB, Bandara R. The use of complementary and alternative medicines (CAMs) in the treatment of diabetes mellitus: is continued use safe and effective? Nutrition journal. 2014;13:102.
24. Parikh NH, Parikh PK, Kothari C. Indigenous plant medicines for health care: treatment of Diabetes mellitus and hyperlipidemia. Chinese journal of natural medicines. 2014;12(5):335-44.
25. Sales DS, Carmona F, de Azevedo BC, Taleb-Contini SH, Bartolomeu AC, Honorato FB, et al. Eugenia punicifolia (Kunth) DC. as an adjuvant treatment for type-2 diabetes mellitus: a non-controlled, pilot study. Phytotherapy research : PTR. 2014;28(12):1816-21.
26. Xin QQ, Liu Y, Yang L, Fu CG, Chen KJ. [Ginkgo preparations of Chinese medicine and treatment of diabetes: mechanisms and clinical applications]. Zhongguo Zhong yao za zhi = Zhongguo zhongyao zazhi = China journal of Chinese materia medica. 2014;39(23):4509-15.
27. Abbas ZK, Saggu S, Sakeran MI, Zidan N, Rehman H, Ansari AA. Phytochemical, antioxidant and mineral composition of hydroalcoholic extract of chicory (Cichorium intybus L.) leaves. Saudi journal of biological sciences. 2015;22(3):322-6.
28. Pushparaj PN, Low HK, Manikandan J, Tan BK, Tan CH. Anti-diabetic effects of Cichorium intybus in streptozotocin-induced diabetic rats. Journal of ethnopharmacology. 2007;111(2):430-4.

29. Ziamajidi N, Khaghani S, Hassanzadeh G, Vardasbi S, Ahmadian S, Nowrouzi A, et al. Amelioration by chicory seed extract of diabetes- and oleic acid-induced non-alcoholic fatty liver disease (NAFLD)/non-alcoholic steatohepatitis (NASH) via modulation of PPARalpha and SREBP-1. *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association*. 2013;58:198-209.
30. Bakhtiary Z. Herbal medicines in diabetes. *Iranian journal of diabetes and obesity*. 2011;3(2):88-95.
31. Carvalho CA, Camargo AM, Cagnon VH, Padovani CR. Effects of experimental diabetes on the structure and ultrastructure of the coagulating gland of C57BL/6J and NOD mice. *Anat Rec A Discov Mol Cell Evol Biol*. 2003;270(2):129-36.
32. Kim M, Shin HK. The water-soluble extract of chicory influences serum and liver lipid concentrations, cecal short-chain fatty acid concentrations and fecal lipid excretion in rats. *The Journal of nutrition*. 1998;128(10):1731-6.
33. Ghamarian A, Abdollahi M, Su X, Amiri A, Ahadi A, Nowrouzi A. Effect of chicory seed extract on glucose tolerance test (GTT) and metabolic profile in early and late stage diabetic rats. *DARU Journal of Pharmaceutical Sciences*. 2012;20(1):56.
34. Tousch D, Lajoix AD, Hosy E, Azay-Milhau J, Ferrare K, Jahannault C, et al. Chicoric acid, a new compound able to enhance insulin release and glucose uptake. *Biochem Biophys Res Commun*. 2008;377(1):131-5.
35. Wight WA, Niekerk JV. Determination of reducing sugars, sucrose and insulin. *Journal of Agriculture and Food Chemistry*. 1983;31:282-5.
36. Norbaek R, Nielsen K, Kondo T. Anthocyanins from flowers of *Cichorium intybus*. *Phytochemistry*. 2002;60(4):357-9.
37. Rees BS, Harborne JB. The role of sesquiterpene lactones and phenolics in the chemical defense of the chicory plant. *Phytochemistry*. 1985;24:2225-31.
38. Lairon D. Dietary fibres: Effects on lipid metabolism and mechanisms of action. *European Journal of Clinical Nutrition*. 1996;50:125-33.