

**Original Article**

Associations between Dietary Total Antioxidant Capacity and Risk of Nonalcoholic Fatty Liver Disease (NAFLD) in Adults: A Case-Control Study

Mohammad Hassan Sohouli ^{1,2*}, Somaye Fatahi ¹

1- Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Tehran, Iran

2- Student Research Committee, Faculty of public health Branch ,Iran University of medical sciences, Tehran, Iran

Received: March 2020

Accepted: May 2020

ABSTRACT

Background and Objectives: Dietary total antioxidant capacity (DTAC) is suggested as a useful tool for the assessment of relationships between cumulative antioxidant food capacities and several chronic disorders. However, relationships between the total antioxidant capacity of the diet (TAC) and the risk of NAFLD has not previously been assessed. The aim of this study was to assess relationships between DTAC and risk of NAFLD in a case-control study.

Materials and Methods: This case-control study was carried out on 158 patients with NAFLD and 357 healthy individuals aged 18–55 years. Dietary data were collected using validated 168-items quantitative food frequency questionnaires. Triglyceride (TG), total cholesterol (TC), high density lipoprotein (HDL-C), low density lipoprotein (LDL-C) and fasting blood glucose (FBS) concentrations were assessed using enzymatic methods and commercial kits. The DTAC was calculated based on the oxygen radical absorbance capacity of each food (except for coffee) reported by US Department of Agriculture. Statistical Analysis was carried out using SPSS Software.

Results: The mean \pm SD (standard deviation) for age and body mass index (BMI) of the study participants were 43.9 years \pm 5.9 and had $30.5 \text{ kg/m}^2 \pm 2.6$. The NAFLD patients included higher BMI and female proportion, compared to control group. The NAFLD patients included higher smoking rates, biochemical parameters (TG, TC, HDL-C, LDL-C and FBS) and DTAC scores, compared to control groups. However, patients with NAFLD included lower HDL levels and physical activities, compared to control group. The highest tertile of DTAC showed lower risks of NAFLD, compared to the lowest tertile. This association was significant after controlling for potential confounders (p for trend <0.001).

Conclusions: Findings suggest that promotion of naturally increased antioxidant capacities may help prevent development of NAFLD.

Keywords: Dietary total antioxidant capacity, Obesity, Nonalcoholic fatty liver disease

Introduction

Nonalcoholic fatty liver disease (NAFLD) is one of the major global health challenges, including several liver abnormalities such as hepatic steatosis, cirrhosis or hepatocellular carcinoma (1). The disease is characterized by accumulation of more than 5% of fats in liver tissue cells in absence of other risk factors, viruses, immune and metabolic disorders and drug abuses (2). Worldwide, estimated prevalence of NAFLD in general populations is nearly 25% (3). In Iran, prevalence of NAFLD in children and adults has been reported as 7 and 35%, respectively (4). Studies have suggested that systemic inflammations or excessive levels of free radicals and disruption of antioxidant balances in the body (generally oxidative stress) may play key roles in pathogenesis of NAFLD

through increased lipid peroxidation in cell membranes (5, 6). When antioxidant and anti-inflammatory defenses are exhausted, chronic states of liver diseases increase. Various aspects of the effects of natural antioxidants in foods on the modulation of oxidative stresses and disorders in antioxidant systems have been studied (7, 8). One of the diet components that can affect these symptoms is the natural antioxidant. Interventional studies using antioxidant supplements (vitamins C and E) have shown that supplementation with antioxidants may include positive effects on these symptoms and diseases (9). However, assessment of an antioxidant compound alone cannot reflect total antioxidant potency of the diet and reflects the synergistic and potential effects of dietary antioxidant

*Address for correspondence: Mohammad Hassan Sohouli, Department of Nutrition, School of Public Health, Iran University of Medical Sciences, Hemmat superhighway, Tehran, Iran. E-mail address: mohammadhassansohouli@gmail.com

interactions. Thus, the term of dietary total antioxidant capacity (DTAC) has been developed as an appropriate tool to assess effects of dietary antioxidants. This includes a strong correlation with serum total antioxidant capacity (10) and is closely linked to the quality of diets to assess risks of chronic diseases (11).

Several evidences have suggested potential links between DTAC and decreased risks of chronic diseases such as diabetes (10), metabolic and oxidative stress markers (11), ulcerative colitis (12), blood pressure (13) and cardiovascular diseases (CVDs) (14), which share common metabolic parameters with NAFLD. However, to the best of the authors' knowledge, associations between DTAC and risks of developmental NAFLD have not been investigated. Only a recent cross-sectional study on steatohepatitis patients has declared that high DTAC score is linked to low hepatic injuries through reducing free radical productions and hence decreasing oxidative stresses (15). Considering lack of convincing evidences on associations of DTAC with liver function, the current study was carried out to investigate possible associations between DTAC and risks of NAFLD in Iranian population.

Materials and Methods

Participants

This case-control study was carried out on 158 patients with NAFLD and 357 healthy individuals referred to Hazrat Rasoul Hospital, Tehran, Iran, 2018–2019. Diagnosis of NAFLD was carried out using chronic elevation of liver enzymes, absence of alcohol consumption, ultrasonography results of the liver showing NAFLD (Grades II and III) and exclusion of other etiologies of liver diseases. Case group patients were newly diagnosed and were not treated before the study. Healthy individuals were considered as control group based on the laboratory tests and liver ultrasonography (not suffering from stages of hepatic steatosis). Control group was selected from other patients referred to other wards of the hospital such as ophthalmology, orthopedics, maxillofacial surgery and ear, nose and throat wards, who were not diagnosed with NAFLD and had no history of alcohol overall or had a history of alcohol consumption less than 10 mg/d in women and less than 20 mg/d in men. Case and control group members were matched regarding age and sex. After entering the study, information on demographic variables and enzyme levels were collected by completing general information Questionnaires. Liver enzyme levels were assessed after return of the participants to the hospital. For the completion of data on dietary intakes and other information, participants were invited to the research center at a special date. Participants with a history of certain diseases (diabetes, CVDs, myocardial infarctions/strokes, cancers, viral hepatitis, Wilson's disease and autoimmune disorders of the liver) were excluded from the list. Pregnant and lactating women and subjects with arbitrary special

diets were excluded from the list as well. In this study, nutritionists were used as interviewers. Therefore, all patients responded completely to the survey questions. Furthermore, written informed consents were completed for the participants. To assess physical activity levels of the participants, general practice physical activity questionnaires (GPPAQ) were used. The GPPAQ is a simple questionnaire reflecting personal current physical activities (16). This study was approved by the Research Council and Ethics Committee, Iran University of Medical Sciences, Tehran, Iran (No. IR.IUMS.REC.1397.667).

Anthropometric assessment

Anthropometric measurements were carried out by a trained dietitian. Weight was measured with minimum clothes and no shoes using standard SECA 700 Digital Scale (SECA, Germany) and recorded to the nearest 100 g. Height was measured in a standing relaxed shoulder position with no shoes to the nearest 0.5 cm using mounted tape (SECA stadiometer, Germany). Body mass index (BMI) was calculated as weight (kg) divided by height in square meters (m^2).

Measurement of biomarkers

After 12 h of fasting, 10 ml of fasting blood were collected between 7 and 10 am from all participants. Blood samples were centrifuged at 500 g for 10 min at 4°C within 30–45 min of the collection. Sera were stored at -80°C until use. Then, triglyceride (TG), total cholesterol (TC), HDL-C, LDL-C and fasting blood glucose (FBS) concentrations were assessed using enzymatic methods and commercial kits (Pars Azmoon, Tehran, Iran), Noor Laboratory, Iran University of Medical Sciences, Tehran, Iran. Alanine aminotransferase (ALT) was assessed using commercially available enzymatic reagents (Pars Azmoon, Tehran, Iran) and auto analyzer (BT-3000). Furthermore, NAFLD was diagnosed using ultrasonography (US), based on echogenicity and visualization of vasculature, parenchyma and diaphragm. These were compared to histological features based on Brunt's Classification (17).

Dietary assessment and DTAC calculation

Trained dietitians administered usual food intakes during interviews. Validated semi-quantitative food frequency questionnaires (FFQ) with 168 food items were used to assess dietary intakes (18). Consumption of food items was calculated on a daily, weekly or monthly basis. Then, data were transformed into average monthly intakes and analyzed using Nutritionist Software v.4. The USDA portion sizes and household measures were used for each food. Moreover, DTAC was calculated based on the oxygen radical absorbance capacity of each food (except for coffee) reported by the US Department of Agriculture and expressed as mmol of trolox equivalent/100 g of food (mmol TE/100 g) (19).

Statistical analysis

Statistical analysis was carried out using Statistical Package Software for Social Science v.21 (SPSS Inc., Chicago, IL, USA). Kolmogorov-Smirnov test and histogram chart were used for testing normality of the data. The independent Student's t-test and Mann-Whitney test were used to compare variables with normal and abnormal distributions, respectively. Baseline characteristics and dietary intakes were expressed as mean \pm SD (standard deviation) or median (25-75 interquartile range) for quantitative variables and number and percentages for qualitative variables. Comparison of data between the two groups was carried out using independent sample t-test and chi-square for continuous and categorical variables, respectively. Using analysis of covariance (ANCOVA), differences of nutrient intakes were compared within DTAC tertiles. A χ^2 test was used for qualitative variables to identify significant differences within quartile categories of DTAC. All confounders (total fat, dietary fiber and CHO) were selected based on the following two strategies of 1) similar literatures, which assessed dietary total antioxidant capacity such as a study by Sotoudeh (2018) (20) that adjusted this index for dietary fiber and fat intake; and 2) confounders of dietary fat and CHO differed between the case and control groups (see Table 2) and included significant associations with exposure (dietary total antioxidant capacity). In general, no studies are available to suggest that these food groups (fat, CHO and fiber) are directly involved in calculation of dietary total anti-oxidant capacity. Binary logistic regression was used to estimate odds ratios (ORs) with 95% confidence intervals (CIs) adjusted for multiple covariates in various models. Data were presented as mean \pm SD and odds ratio with 95% confidence intervals. Significance levels were reported when $p<0.05$.

Results

The mean \pm SD for age and BMI of the participants were 43.9 years \pm 5.9 (females=58.7%) and 30.5 kg/m 2 \pm 2.6,

respectively. Table 1 demonstrates anthropometric and biochemical indices within case (non-alcoholic fatty liver patients) and control groups. No significant differences were seen between the case and control groups in age and sex ($p=0.76$) and ($p=0.51$), respectively. However, NAFLD patients included higher BMI, compared to control group ($p < 0.001$). Moreover, NAFLD cases include higher levels of smoking, FBS, TG, LDL, TC and ALT, compared to control group ($p < 0.05$). However, patients with NAFLD included lower HDL and DTAC scores as well as physical activities, compared to control group ($p < 0.05$). Dietary intakes in NAFLD case and control group are presented in Table 2.

Significant differences were demonstrated in intakes of more micronutrients (calcium, vitamin E, vitamin B₁₂, vitamin D and vitamin C) and macronutrients (fats, saturated fatty acids, carbohydrates, fruits, vegetables, whole grains, refined grains and fructose) between the case and control groups ($p<0.05$). Additionally, no significant differences were found between the case and control groups in intakes of proteins, fibers, selenium and folates. Participants' characteristics and biochemical parameters within the tertiles of DTAC are shown in Table 3. The mean age significantly increased within the tertiles of DTAC ($p<0.002$), while FBS, TG and ALT significantly decreased within the tertiles of DTAC ($p<0.05$). No significant differences were seen in other characteristics and biochemical parameters and BMI within the tertiles of DTAC. The ORs and 95% CIs for NAFLD in the tertiles of DTAC are described in Table 4. Compared to participants in the lowest tertile of DTAC, those in the highest tertile included significantly lower ORs for NAFLD (crude model: OR, 0.37; 95% CI, 0.24-0.58; p for trend < 0.001), which were still significant after further adjustments for BMI, physical activities and dietary intakes of fats, fibers, carbohydrates and energy as well as biochemical parameters (Model 3: OR, 0.22; 95% CI, 0.11-0.43; p for trend <0.001).

Table 1. Anthropometric and biochemical parameters of the case (nonalcoholic fatty liver patients) and control groups

Variable	Group, mean \pm SD		<i>P</i> value
	Case (n=158)	Control (n=357)	
Women, % (No)	46.3 (98)	45.9 (88)	0.51 ^a
Age, y	45.75 \pm 5.9	45.13 \pm 5.9	0.76 ^b
BMI, kg/m ²	33.19 \pm 3.1	27.95 \pm 2.1	<0.001 ^b
Physical Activity (Met.min/wk)	984.5 \pm 1241.5	1563.2 \pm 1216.7	<0.001 ^b
Smoking (yes), n (%)	16 (7.1)	12 (2.7)	0.006
FBS, mg/dl	109.29 \pm 7.6	92.21 \pm 6.7	<0.001 ^b
Total cholesterol, mg/dl	184.79 \pm 42.1	182.85 \pm 40.1	<0.001 ^b
Triglyceride, mg/dl	180.40 \pm 14.3	130.99 \pm 12.2	0.001 ^b
HDL, mg/dl	41.26 \pm 15.1	48.5 \pm 17.3	0.017 ^b
LDL, mg/dl	121.17 \pm 23.4	109.14 \pm 13.0	<0.001 ^b
ALT, mg/dl	58.50 \pm 24.1	20.53 \pm 13.01	<0.001 ^b
DTAC, mmol TE/100 g	12323.6 \pm 5398.5	17563.4 \pm 8247.2	<0.001 ^b

^a P values are resulted from chi square

^b P values are resulted from student t-test

Table 2. Dietary intakes by the case (nonalcoholic fatty liver patients) and control groups

Variable	Group, mean \pm SD		<i>P</i> value ^a
	Case (n=158)	Control (n=357)	
Energy, kcal	2741.8 \pm 819.89	2427.67 \pm 798.09	0.04
Macronutrients			
Protein, gr	97.33 \pm 34.37	104.47 \pm 39.61	0.06
Fat, gr	103.44 \pm 40.11	86.27 \pm 32.58	<0.001
Saturated fatty acid	32.13 \pm 6.35	26.87 \pm 6.42	0.006
Carbohydrate, gr	408.06 \pm 148.95	378.44 \pm 112.63	0.03
Fiber, gr	46.36 \pm 18.25	47.58 \pm 19.76	0.5
Fruits	249.38 \pm 190.29	368.03 \pm 192.15	<0.001
vegetables	269.36 \pm 183.63	339.31 \pm 185.25	<0.001
Whole grain	121.54 \pm 80.71	134.44 \pm 79.91	<0.001
Refined grain	420.41 \pm 140.35	368.87 \pm 141.65	<0.001
fructose	25.95 \pm 12.19	23.52 \pm 10.33	0.03
<i>Micronutrients</i>			
Calcium (mg/d)	1125.66 \pm 381.61	1311.17 \pm 441.93	<0.001
Selenium(mg/d)	135.39 \pm 56.28	133.34 \pm 52.41	0.717
Vitamin E (mg/d)	13.22 \pm 5.866	18.98 \pm 7.146	<0.001
Folate (mcg/d)	615.53 \pm 182.21	642.29 \pm 194.52	0.176
Vitamin B12 (mcg/d)	6.37 \pm 4.47	5.36 \pm 4.42	0.029
Vitamin D (mcg/d)	1.56 \pm 1.27	2.03 \pm 1.80	0.03
Vitamin C (mg/d)	175.72 \pm 88.09	231.92 \pm 144.06	<0.001

^aP values are resulted from student t-test

Table 3. Characteristics and biochemical parameters within tertiles of dietary total antioxidant capacity in the study population

	Tertiles of dietary total antioxidant capacity			<i>P</i> ^a
	T1	T2	T3	
Age(year)	37.0 \pm 8.3	38.2 \pm 8.5	39.5 \pm 9.6	0.002
BMI, kg/m ²	26.8 \pm 4.1	26.7 \pm 4.4	26.9 \pm 4.4	0.687
Physical Activity (Met.min/wk)	1462 \pm 862	1454 \pm 850	1368 \pm 939	0.261
Smoking (yes), n (%)	9 (3.5)	11 (4.9)	8 (4.2)	0.722
FBS, mg/dl	102.73 \pm 43.92	95.20 \pm 36.79	94.81 \pm 33.58	0.001
Total cholesterol, mg/dl	186.34 \pm 41.68	182.20 \pm 43.70	183.27 \pm 46.52	0.804
Triglyceride, mg/dl	155.56 \pm 114.17	144.46 \pm 72.26	141.09 \pm 79.87	<0.001
HDL, mg/dl	47.09 \pm 12.02	44.34 \pm 12.00	47.52 \pm 14.61	0.198
LDL, mg/dl	113.42 \pm 31.53	111.98 \pm 35.75	111.11 \pm 35.77	0.261
ALT, mg/dl	39.24 \pm 51.22	30.49 \pm 25.78	29.78 \pm 38.03	<0.001

^aP-value from one-factor ANCOVA test or χ^2 test, for continuous or categorical variables, respectively.

Table 4. Associations between tertiles of dietary total antioxidant capacity and risks of NAFLD in participants

	Tertile of dietary total antioxidant capacity			<i>P</i> for trend
	T1	T2	T3	
DTAC				
Case/Total	61 / 105	58 / 125	39/ 127	
Crude	1.00 (Ref)	0.76(0.52 - 1.12)	0.37 (0.24 - 0.58)	< 0.001
Model 1*	1.00 (Ref)	0.74 (0.42 - 1.28)	0.20 (0.10 - 0.40)	< 0.001
Model 2†	1.00 (Ref)	0.93 (0.48 - 1.77)	0.29 (0.14 - 0.61)	0.001
Model 3‡	1.00 (Ref)	0.68 (0.39 - 1.19)	0.22 (0.11 - 0.43)	< 0.001

Binary logistic regression was used to estimate odds ratios (ORs) and 95% confidence intervals (CIs) adjusted for multiple covariates in different model.

* Model 1: adjusted for age and sex

† Model 2: adjusted for model 1 and BMI, physical activity, dietary intake of fat, fiber, carbohydrate and energy.

‡ Model 3: additionally adjusted for fasting blood sugar, TG, cholesterol, LDL-C, and HDL-C at baseline.

Discussion

The current case-control study investigated associations between DTAC and NAFLD. Generally, DTAC included inverse associations with weight, FBS, TG and ALT. Participants with higher DTAC scores included significantly lower ORs in NAFLD, after adjustment for BMI, physical activity, education level, and dietary intakes of fibers, fats, energy, coffee as well as biochemical parameters, compared to low DTAC scores. In the present study, case group included lower DTAC scores, compared to control group, suggesting that DTAC may be a potential predictor of the risks of developing NAFLD. To the best of the authors' knowledge, the current study is the first documentation of inverse associations between DTAC and risks of NAFLD. In previous studies, dietary TAC has inversely been associated with several chronic diseases such as CVDs (14), cancers (21), diabetes (10) and metabolic disorders (11), which share common metabolic parameters with NAFLD. Only one case-control study has assessed status of the blood redox with the severity of NAFLD (22) with no differences in dietary antioxidant scores between the two groups of case and control. In the highlighted study, no calculations of the odds or risks of the disease were carried out and the sample size was small. Although, thiobarbituric acid reactive substances were significantly associated with the likelihood of NAFLD, independently diet's total antioxidant.

Overall, inverse associations were reported between intakes of antioxidant micronutrients and antioxidant-rich foods and risks of NAFLD. For example, DASH diets (dietary approaches to stop hypertension), dietary patterns with antioxidant-rich components based on fruits and vegetables, low-fat dairy products and whole grains demonstrated inverse relationships with risks of NAFLD (23, 24) as well as inverse associations in chronic diseases associated with NAFLD such as CVDs and diabetes (25). Furthermore, coffee has been reported as an antioxidant-rich nutrient for the risk reduction of NAFLD. Recently, a meta-analysis has suggested that coffee intake in a dose-dependent manner decreases risks of developing NAFLD (26). Ability of other antioxidant-rich nutrients to increase plasma TAC and hence decrease risks of NAFLD through modifications of oxidative stress has been well documented. These nutrients include bayberry juice (27), chocolate (28), onion (29), lettuce (30) and tomato products (31). Findings have been verified even in supplementary interventions with antioxidants. Furthermore, beneficial effects of the combination of antioxidant supplements have been reported (32), as synergistic effects of simultaneous administration and combination of one or more antioxidant supplements decrease risks of NAFLD (32).

Oxidative stress has been shown to involve in pathogenesis and development of lipid metabolism

disorders and insulin resistance and can lead to NAFLD (33). Decreases in antioxidant defense mechanisms can increase lipid peroxidation, damage cellular organs and enzymes and cause insulin resistance (34). Thus, high dietary antioxidants can improve lipid and glucose metabolism disorders and decrease risks of NAFLD by protecting liver cells (8, 15, 34). Additionally, high intakes of antioxidant-rich nutrients from plant foods, as part of a healthy diets, may include health benefits not only by protecting cells from oxidative damages (35), but also by providing fibers and antioxidant nutrients such as vitamin D, vitamin C, vitamin E and magnesium, which include advantageous for BMI, serum lipids and glucose (36). Beneficial synergistic effects of antioxidants, fibers, potassium, magnesium and other phytochemicals on prevention of NAFLD have been observed in various studies (36, 37). In contrast, overweight and abdominal obesity have been suggested as potential risk factors for the progression of NAFLD (38).

Dietary antioxidants provide protective mechanisms against obesity linked disorders, including inhibition of fat absorption, promotion of catabolism in adipose tissues, inhibition of proliferation, differentiation and angiogenesis in preadipocytes and induction of apoptosis in mature adipocytes (39). Regarding these mechanisms, the mean weight of patients significantly decreased within the tertiles of DTAC. However, the present study included limitations that might affect interpretation of the results. Similar to other case-control studies, causal relationships between DTAC and NAFLD were not investigated. Moreover, use of 168-item FFQ questionnaires caused tired participants and biases in their responses, which were resolved by a trained questioner. Furthermore, conditions of foods such as growth, cultivation, storage, processing and cooking conditions and the assay methods might affect antioxidant contents of the foods (19). Despite these limitations, this was the first study to investigate relationships between DTAC and NAFLD using a case control design. However, effects of confounders were tried to eliminate as far as possible using adjustment of a wide range of variables and validated questionnaires.

Conclusion

Findings have shown that high DTACs are associated with decreased risks of NAFLD in adults, suggesting that promotion of naturally elevated antioxidant capacities may help prevent developments of NAFLD. Encouraging consumption of diets with high antioxidant capacities is important in nutritional interventions for the prevention of NAFLD. However, complementary studies are necessary to further investigate associations of DTAC intakes with risks of NAFLD in other settings.

Acknowledgement

The authors appreciate participants of this study.

Financial disclosure

The authors declare no personal or financial conflicts.

Authors' contributions

S. F. and M. S. contributed to conception, design and statistical analysis. M. S. and N. H. contributed to data collection and manuscript drafting. F. S. and A. S. supervised the study. All authors approved the final version of the manuscript.

References

1. Petaja EM, Yki-Jarvinen H. Definitions of normal liver fat and the association of insulin sensitivity with acquired and genetic NAFLD—a systematic review. *International journal of molecular sciences*. 2016;17(5):633.
2. Than NN, Newsome PN. A concise review of non-alcoholic fatty liver disease. *Atherosclerosis*. 2015;239(1):192-202.
3. Anderson EL, Howe LD, Jones HE, Higgins JP, Lawlor DA, Fraser A. The prevalence of non-alcoholic fatty liver disease in children and adolescents: a systematic review and meta-analysis. *PLoS one*. 2015;10(10).
4. Moghaddasifar I, Lankarani K, Moosazadeh M, Afshari M, Ghaemi A, Aliramezany M, et al. Prevalence of non-alcoholic fatty liver disease and its related factors in Iran. *International journal of organ transplantation medicine*. 2016;7(3):149.
5. Ashraf N, Sheikh T. Endoplasmic reticulum stress and oxidative stress in the pathogenesis of non-alcoholic fatty liver disease. *Free radical research*. 2015;49(12):1405-18.
6. Cichoñ-Lach H, Michalak A. Oxidative stress as a crucial factor in liver diseases. *World journal of gastroenterology: WJG*. 2014;20(25):8082.
7. Serafini M, Del Rio D. Understanding the association between dietary antioxidants, redox status and disease: is the total antioxidant capacity the right tool? *Redox report*. 2004;9(3):145-52.
8. Meydani M. Dietary antioxidants modulation of aging and immune-endothelial cell interaction. *Mechanisms of ageing and development*. 1999;111(2-3):123-32.
9. Ivancovsky-Wajcman D, Fliss-Isakov N, Salomone F, Webb M, Shibolet O, Kariv R, et al. Dietary vitamin E and C intake is inversely associated with the severity of nonalcoholic fatty liver disease. *Digestive and Liver Disease*. 2019;51(12):1698-705.
10. Mancini FR, Affret A, Dow C, Balkau B, Bonnet F, Boutron-Ruault M-C, et al. Dietary antioxidant capacity and risk of type 2 diabetes in the large prospective E3N-EPIC cohort. *Diabetologia*. 2018;61(2):308-16.
11. Hermsdorff HHM, Puchau B, Volp ACP, Barbosa KB, Bressan J, Zulet MÁ, et al. Dietary total antioxidant capacity is inversely related to central adiposity as well as to metabolic and oxidative stress markers in healthy young adults. *Nutrition & metabolism*. 2011;8(1):59.
12. Koch TR, Yuan L-X, Stryker SJ, Ratliff P, Telford GL, Opara EC. Total antioxidant capacity of colon in patients with chronic ulcerative colitis. *Digestive diseases and sciences*. 2000;45(9):1814-9.
13. Villaverde P, Lajous M, MacDonald C-J, Fagherazzi G, Bonnet F, Boutron-Ruault M-C. High dietary total antioxidant capacity is associated with a reduced risk of hypertension in French women. *Nutrition journal*. 2019;18(1):31.
14. Rautiainen S, Levitan EB, Orsini N, Åkesson A, Morgenstern R, Mittleman MA, et al. Total antioxidant capacity from diet and risk of myocardial infarction: a prospective cohort of women. *The American journal of medicine*. 2012;125(10):974-80.
15. de Oliveira DG, de Faria Ghetti F, Moreira APB, Hermsdorff HHM, de Oliveira JM, de Castro LEVV. Association between dietary total antioxidant capacity and hepatocellular ballooning in nonalcoholic steatohepatitis: a cross-sectional study. *European journal of nutrition*. 2019;58(6):2263-70.
16. Ahmad S, Harris T, Limb E, Kerry S, Victor C, Ekelund U, et al. Evaluation of reliability and validity of the General Practice Physical Activity Questionnaire (GPPAQ) in 60-74 year old primary care patients. *2015;16(1):113*.
17. Shannon A, Alkhouri N, Carter-Kent C, Monti L, Devito R, Lopez R, et al. Ultrasonographic quantitative estimation of hepatic steatosis in children with nonalcoholic fatty liver disease (NAFLD). *Journal of pediatric gastroenterology and nutrition*. 2011;53(2):190.
18. Azadbakht L, Esmaillzadeh A. Red meat intake is associated with metabolic syndrome and the plasma C-reactive protein concentration in women. *The Journal of nutrition*. 2009;139(2):335-9.
19. Haytowitz DB, Bhagwat S. USDA database for the oxygen radical absorbance capacity (ORAC) of selected foods, Release 2. US Department of Agriculture. 2010:10-48.
20. Sotoudeh G, Abshirini M, Bagheri F, Siassi F, Koohdani F, Aslany Z. Higher dietary total antioxidant capacity is inversely related to prediabetes: a case-control study. *Nutrition*. 2018;46:20-5.
21. Vece MM, Agnoli C, Grioni S, Sieri S, Pala V, Pellegrini N, et al. Dietary total antioxidant capacity and colorectal cancer in the Italian EPIC cohort. *PLoS One*. 2015;10(11).
22. Georgoulis M, Fragopoulou E, Kontogianni MD, Margariti A, Boulamatsi O, Detopoulou P, et al. Blood redox status is associated with the likelihood of nonalcoholic fatty liver disease irrespectively of diet's total antioxidant capacity. *Nutrition Research*. 2015;35(1):41-8.
23. Hekmatdoost A, Shamsipour A, Meibodi M, Gheibizadeh N, Eslamparast T, Poustchi H. Adherence to the dietary approaches to stop hypertension (DASH) and risk of nonalcoholic fatty liver disease. *International journal of food sciences and nutrition*. 2016;67(8):1024-9.
24. Xiao M-L, Lin J-S, Li Y-H, Liu M, Deng Y-Y, Wang C-Y, et al. Adherence to the Dietary Approaches to Stop Hypertension (DASH) diet is associated with lower presence of non-alcoholic fatty liver disease in middle-aged and elderly adults. *Public Health Nutrition*. 2019;1-9.
25. Liese AD, Nichols M, Sun X, D'Agostino RB, Haffner SM. Adherence to the DASH Diet is inversely associated with incidence of type 2 diabetes: the insulin resistance atherosclerosis study. *Diabetes care*. 2009;32(8):1434-6.
26. Chen Y-P, Lu F-B, Hu Y-B, Xu L-M, Zheng M-H, Hu E-D. A systematic review and a dose-response meta-analysis of

coffee dose and nonalcoholic fatty liver disease. *Clinical Nutrition*. 2019;38(6):2552-7.

27. Guo H, Zhong R, Liu Y, Jiang X, Tang X, Li Z, et al. Effects of bayberry juice on inflammatory and apoptotic markers in young adults with features of non-alcoholic fatty liver disease. *Nutrition*. 2014;30(2):198-203.

28. Malhi H, Loomba R. dark chocolate may improve NAFLD and metabolic syndrome by reducing oxidative stress. *Alimentary pharmacology & therapeutics*. 2016;44(5):533.

29. Emamat H, Foroughi F, Eini-Zinab H, Taghizadeh M, Rismanchi M, Hekmatdoost A. The effects of onion consumption on treatment of metabolic, histologic, and inflammatory features of nonalcoholic fatty liver disease. *Journal of Diabetes & Metabolic Disorders*. 2015;15(1):25.

30. Nikkhah-Bodaghi M, Ghanavati M, Hekmatdoost A. Polyphenol intakes and risk of impaired lipid profile, elevated hepatic enzymes and nonalcoholic fatty liver disease. *Nutrition & Food Science*. 2019.

31. García-Alonso F, González-Barrio R, Martín-Pozuelo G, Hidalgo N, Navarro-González I, Masuero D, et al. A study of the prebiotic-like effects of tomato juice consumption in rats with diet-induced non-alcoholic fatty liver disease (NAFLD). *Food & function*. 2017;8(10):3542-52.

32. Sawangjit R, Chongmelaxme B, Phisalprapa P, Saokaew S, Thakkinstian A, Kowdley KV, et al. Comparative efficacy of interventions on nonalcoholic fatty liver disease (NAFLD): A PRISMA-compliant systematic review and network meta-analysis. *Medicine*. 2016;95(32).

33. Narasimhan S, Gokulakrishnan K, Sampathkumar R, Farooq S, Ravikumar R, Mohan V, et al. Oxidative stress is independently associated with non-alcoholic fatty liver disease (NAFLD) in subjects with and without type 2 diabetes. *Clinical biochemistry*. 2010;43(10-11):815-21.

34. Liu W, Baker SS, D Baker R, Zhu L. Antioxidant mechanisms in nonalcoholic fatty liver disease. *Current drug targets*. 2015;16(12):1301-14.

35. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *The international journal of biochemistry & cell biology*. 2007;39(1):44-84.

36. Mouzaki M, Allard JP. The role of nutrients in the development, progression, and treatment of nonalcoholic fatty liver disease. *Journal of clinical gastroenterology*. 2012;46(6):457-67.

37. Dreher ML. Dietary Patterns, Foods, Nutrients and Phytochemicals in Non-Alcoholic Fatty Liver Disease. *Dietary Patterns and Whole Plant Foods in Aging and Disease*: Springer; 2018. p. 291-311.

38. Jakobsen MU, Berentzen T, Sørensen T, Overvad K. Abdominal obesity and fatty liver. *Epidemiologic reviews*. 2007;29(1):77-87.

39. Barnard ND, Scialli AR, Turner-McGrievy G, Lanou AJ, Glass J. The effects of a low-fat, plant-based dietary intervention on body weight, metabolism, and insulin sensitivity. *The American journal of medicine*. 2005;118(9):991-7.