**Original Article**

## Improving Oxidative Stability of Virgin Olive Oil: Comparison of *Zataria Multiflora* Essential Oil with $\alpha$ -Tocopherol

Malihe Keramat<sup>1</sup>, Mohammad-Taghi Golmakani\*<sup>2</sup>, Mahmoud Aminlari<sup>3</sup>, Seyed Shahram Shekarforoush<sup>4</sup>

1- PhD Student, Department of Food Science and Technology, Faculty of Agriculture, Shiraz University, Shiraz, Iran

2- Associate Professor, Department of Food Science and Technology, School of Agriculture, Shiraz University, Shiraz, Iran

3- Professor, Department of Basic Sciences, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

4- Professor, Department of Hygiene and Food Quality Control, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

Received: December 2017

Accepted: February 2018

### ABSTRACT

**Background and Objectives:** Virgin olive oil is a vastly consumed product, with widespread appreciation for its good nutritional and health properties. However, oxidation can reduce its quality. The aim of this study was to investigate how the essential oil of *Zataria multiflora* (Shirazi thyme) can contribute to the prevention of virgin olive oil oxidation in comparison with the actions of  $\alpha$ -tocopherol and BHT. Furthermore, the synergistic activities of citric acid with BHT, *Z. multiflora* essential oil, and  $\alpha$ -tocopherol were investigated.

**Materials and Methods:** Antioxidant activity of the essential oil was determined using radical scavenging capacity and reducing power assays. Virgin olive oil samples were stored at  $60\pm 1$  °C in closed amber bottles for 16 days. Oxidation levels of samples were determined by measuring peroxide, anisidine, TOTOX,  $K_{232}$ ,  $K_{268}$  values, and chlorophyll and carotenoid contents of the samples during the storage period.

**Results:** *Z. multiflora* essential oil exhibited a significant radical scavenging capacity and reducing power. Peroxide, anisidine, TOTOX,  $K_{232}$ , and  $K_{268}$  values of samples containing *Z. multiflora* essential oil were significantly lower than those of the control group (without antioxidants). *Z. multiflora* essential oil reduced the oxidation of virgin olive oil to the same extent as BHT did. *Z. multiflora* essential oil was more effective than  $\alpha$ -tocopherol. The synergistic activities between citric acid and the various compounds, i.e. BHT, *Z. multiflora* essential oil, and  $\alpha$ -tocopherol were 2.42%, 4.74%, 1.28% respectively.

**Conclusions:** In general, *Z. multiflora* essential oil can be considered as natural antioxidant for the stabilization of virgin olive oil against oxidation.

**Keywords:** Accelerated storage, Oxidation, Virgin olive oil, *Zataria multiflora*

### Introduction

Virgin olive oil (VOO) is commonly accepted by consumers as an oil produced without any chemical treatments. It has a high oxidative stability, mainly because of two reasons; firstly, it has a high ratio of monounsaturated-polyunsaturated fatty acids and, secondly, it has significant amounts of polyphenol compounds with strong antioxidant activities (1). However, it contains polyunsaturated fatty acids such as linoleic acid and linolenic acid which reduce VOO oxidative stability. The most important cause of VOO quality degradation is oxidation. Oxidation occurs by enzymatic and chemical reaction pathways (2).

In order to improve the stability of fats and oils, synthetic antioxidants, such as butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), and tert-butyl hydroquinone (TBHQ) are added into fats and oils (3). However, it has been shown that BHA and BHT are not stable at high temperatures (4,5). Furthermore, TBHQ is not allowed for food application in Canada, Japan, and Europe. In addition, BHA has been removed from the list of generally recognized as safe (GRAS) compounds (6). These health concerns have increased the demand for natural antioxidants (7). Herbs and

\*Address for correspondence: Mohammad-Taghi Golmakani, Associate Professor, Department of Food Science and Technology, School of Agriculture, Shiraz University, Shiraz, Iran. E-mail: golmakani@shirazu.ac.ir

spices contain high amounts of biological active compounds with significant antioxidant activity (3).

*Zataria multiflora* belongs to the Lamiaceae family. This aromatic plant is known as Avishan Shirazi. *Z. multiflora* is native to Iran, Pakistan, and Afghanistan (8). The main components of *Z. multiflora* essential oil (EO) are oxygenated monoterpenes thymol and carvacrol, and their corresponding monoterpene hydrocarbon precursors, i.e.  $\gamma$ -terpinene and *p*-cymene. (9). It has been shown that thymol and carvacrol can pose high antioxidant activities. Thymol and carvacrol can donate hydrogen atoms from their phenol hydroxyl groups to peroxy radicals and terminate lipid peroxidation chain reactions (10, 11).

Many researches have investigated the effectiveness of different herbs and spices on the stability of vegetable oils. It has been observed that Ajwain EO (0.075%) lowers the oxidation of sunflower oil to a higher extent than BHA and BHT (0.02%) did (12). Oregano EO was effective against extra VOO oxidation (13). Also, there is considerable evidence that shows the synergistic effects of various antioxidants in vegetable oils. Hras *et al.* reported that rosemary extract showed synergistic effects with ascorbyl palmitate and CA in sunflower oil (14). Also, it has been reported that a combined mixture of honey and mint exhibited a synergistic activity and could be used as a promising food additive (15).

The goal of this study was to investigate the effect of *Z. multiflora* EO on VOO oxidation in comparison with  $\alpha$ -tocopherol and BHT. Also, the synergistic effects of CA with *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were investigated.

## Materials and Methods

**Materials:** Dried aerial parts of *Z. multiflora* were purchased from a local market in Shiraz, Iran. The genus and species of *Z. multiflora* was authenticated by taxonomists from the Herbarium of Biology Department at Shiraz University, Shiraz, Iran. The VOO was provided by Etko Oil organization. Chemicals such as  $\alpha$ -tocopherol, 2, 2-diphenyl-1-picrylhydrazyl radical (DPPH<sup>•</sup>), BHT, TBHQ, potassium ferricyanide, sodium phosphate, trichloroacetic acid, iron (III) chloride, copper (II) chloride, neocuproine ( $\geq 98\%$ ), ammonium acetate, and Folin–Ciocalteu reagent were purchased from Sigma-Aldrich (St. Louis, MO). Potassium iodide, cyclohexane, methanol, isooctane, hexane, and

methanol were purchased from Merck (Darmstadt, Germany).

**Extraction of EO:** *Z. multiflora* EO was extracted according to the method described by Golmakani and Rezaei (9) using a Clevenger-type apparatus. Briefly, 50 grams of *Z. multiflora* sample was mixed with 500 mL of distilled water and hydrodistilled for 3 h. EO samples were dried over anhydrous sodium sulphate and were stored at -18 °C until further analysis (9).

**Gas Chromatography (GC) Analysis of EO:** GC analysis of the EO was carried out using the method described by Khajehie *et al.* (16). The EO constituents were identified using a GC (7890A, Agilent Technologies, Santa Clara, CA) which was coupled with a mass spectrometer (5975C, Agilent Technologies, Santa Clara, CA) operating at 70 eV ionization energy, 0.5 s/scan and a mass range of 35-400 atomic mass units (amu). HP-5MS capillary column (5% Phenyl Polysilphenylene-siloxane, 30 m length, 0.25 mm internal diameter, and 0.25  $\mu$ m film thickness, Agilent Technologies, Santa Clara, CA) was used. The carrier gas was Helium at a flow rate of 0.9 mL/min. In addition, quantitative determination of EO constituents was carried out under the same chromatographic conditions using a GC equipped with a flame ionization detector (FID). For each compound, the relative data for percentage was determined from the electronic integration of the chromatogram's peak areas.

## Antioxidant Activity of EO

**Free Radical Scavenging Capacity:** The free radical scavenging capacities of BHT and *Z. multiflora* EO were determined according to the method of Farahmand *et al.* (17), using the DPPH<sup>•</sup> assay. The results were expressed as IC<sub>50</sub> value, which is defined as the amount of sample concentration that can scavenge 50% of DPPH<sup>•</sup>.

**Reducing Assay:** The ferric reducing antioxidant power (FRAP) of the EO was measured using the procedure described by Farahmand *et al.* (17), by reducing the Fe (III) to Fe (II). The cupric ion reducing antioxidant capacity (CUPRAC) of the EO was measured by reducing the Cu (II) to Cu (I), using the method of Apak *et al.* (18). In both FRAP and CUPRAC assays, vitamin C was used as the positive control. FRAP and CUPRAC results were expressed as mg of vitamin C equivalents per gram of sample (mg VCE/g).

## Initial Quality of VOO

**Free Fatty Acid Content:** Free fatty acids were measured according to the AOCS official method (Cd 3d-63). The free fatty acid content was calculated as the percentage of oleic acid (19).

**Fatty Acid Analysis:** Fatty acid analysis was carried out using the method of Golmakani *et al.* with some modifications (20). In order to prepare fatty acid methyl esters, VOO was mixed with methanol: acetyl chloride solution at the ratio of 95:5 v/v. The mixture was sealed in a PTFE-lined vial and heated at 85 °C for 1 h. The vial was cooled and 5.0 mL double distilled water was added. Afterwards, 1.0 mL of hexane containing 0.01% TBHQ was added to the mixture. Finally, 1.0 mL of hexane containing 0.01% TBHQ was added again and centrifuged at 4000 ×g at 25 °C for 5 min. The upper phase (hexane layer) obtained after centrifugation contained mixtures of fatty acid methyl esters.

To determine fatty acid composition of VOO, a GC (3420A, Beifen system, China) equipped with a FID was used. A BPX70 capillary column (Bis-cyanopropylsiloxane-silphenylene, 30 m × 0.25 mm internal diameter with 0.25 μm film thickness) was used for the separation of fatty acid methyl esters. The column initial temperature was held at 140 ° for 5 min. After that, it was increased to 180 °C at a rate of 20 °C/min. It was held at 180 °C for 9 min. finally, it was increased to 200 °C at a rate of 20 °C/min and held at that temperature for 3 min. The injector and detector temperatures were set at 250 and 300 °C, respectively. One μL of fatty acid methyl ester samples was injected in to the GC system in split mode (at 1:10 ratio). The carrier gas was nitrogen.

**Calculation of Iodine Value:** Iodine value (IV) was calculated according to the equation described by Kyriakidis and Katsiloulis according to eq. (1).

$$IV = xC1 + yC2 + zC3; x = 0.95, y = 1.6, \text{ and } z = 2.62. \quad \text{eq. (1)}$$

C1, C2, and C3 correspond to the sum of mono-, di-, and tri-unsaturated fatty acid methyl esters, respectively (21).

**Total Phenolic Content:** Total phenolic content of VOO was determined using the Folin-Ciocalteu assay according to the method of Casal *et al.* (22). Results were expressed as μg of gallic acid per g of VOO (μg GAE/g).

**Oxidation Indices of VOO:** The peroxide value (PV) was measured using the AOCS official method

(Cd 8-53). PV results were expressed as milli-equivalents of active oxygen per kg of VOO (19). The anisidine value (AV) was measured using the procedure described by the AOCS official method (Cd 8-53). AV results were expressed as mg per kg of VOO (19). The TOTOX value was calculated using eq. (1) (10).

$$\text{TOTOX value} = AV + 2PV \quad \text{eq. (1)}$$

The  $K_{232}$  and  $K_{268}$  extinction coefficients were measured following the AOCS official method (ch 5-91) (18). Chlorophyll and carotenoid contents were determined using the procedure described by Minguéz-Mosquera *et al.* (23).

**Oxidative Stability of VOO during Accelerated Storage:** *Z. multiflora* EO,  $\alpha$ -tocopherol, and BHT were added to the VOO in concentrations of 1000, 100, and 100 ppm, respectively. Also, mixtures of the EO,  $\alpha$ -tocopherol, and BHT with 100 ppm CA were added to the VOO. VOO samples were placed into amber bottles (60 mL). The bottles were completely filled and sealed. No headspace was left in the bottles. The samples were kept in an incubator at 60±1 °C for 16 days. The oxidation level of VOO samples was determined by measuring PV, AV, TV,  $K_{232}$ , and  $K_{268}$  values every 4 days. Also, chlorophyll and carotenoid contents were determined at the beginning and at the end of the storage period.

The IP is defined as the number of days taken for a sample to reach a PV of 20 meq O<sub>2</sub>/kg (11).

Protection factor (PF) was calculated using eq. (2).

$$PF = \frac{IP_a}{IP_v} \quad \text{eq. (2)}$$

where  $IP_v$  is the induction period (IP) of the VOO sample without antioxidant (control) and  $IP_a$  is IP of the VOO samples containing antioxidant (BHT,  $\alpha$ -tocopherol, and *Z. multiflora* EOs) (13). The effectiveness of antioxidants can be determined using the following scale for the PF values: 1.0-1.5 (very low), 1.5-2.0 (low), 2.0-2.5 (medium), 2.5-3.0 (high), and >3.0 (very high) (24).

Antioxidant activity (AA) functions by the concentration of the antioxidant. AA was determined following eq. (3).

$$AA = \frac{IP_a - IP_v}{[AH] IP_v} \quad \text{eq. (3)}$$

Where [AH] is the antioxidant concentration in ppm unit (25).

Also, the synergistic activity was calculated according to eq. (4).

eq. (4)

$$\text{Synergism (\%)} = \left( \frac{(IP_c - IP_v) - (IP_1 - IP_v) - (IP_2 - IP_v)}{(IP_c - IP_v)} \right) \times 100$$

Where  $IP_v$  is the induction period of the sample treated with the mixture of the antioxidant with CA, the  $IP_1$  is the induction period of the sample treated with the antioxidant without CA, and  $IP_2$  is the induction period of the sample treated with CA (14).

**Statistical Analysis:** All results are reported as the mean value  $\pm$  standard deviation of triplicate measurements. In order to compare the mean values, a general linear model (GLM) procedure from SAS (Statistical Analysis Software, version 9.1; SAS Institute Inc. Cary, NC) was used. Microsoft Office Excel 2010 was used to calculate the regression equations of the oxidation indices from the storage study of VOO.

## Results

**Analysis of EO:** The chemical composition of *Z. multiflora* EOs is shown in Table 1. *Z. multiflora* EO contained high amounts of oxygenated monoterpenes and made up 69.49% of the EO. The monoterpene hydrocarbons made up 27.46% of the *Z. multiflora* EO, while sesquiterpene hydrocarbons comprised 2.97%. The main components of *Z. multiflora* EO were thymol (52.80%), *p*-cymene (11.74%), carvacrol (10.33%), and  $\gamma$ -terpinene (6.06%).

**Antioxidant Activity of EO:** *Z. multiflora* EO was able to scavenge DPPH $^\circ$  with  $IC_{50}$  value of 0.06 mg/ml. Comparison of the DPPH $^\circ$  scavenging capacity of the investigated EO and those expressed by BHT ( $IC_{50}$  = 0.02 mg/ml) showed that the EO possessed slightly lower scavenging activities than BHT. Ferric reducing power and cupric ion reducing power of *Z. multiflora* EO were  $815.96 \pm 86.44$  and  $895.43 \pm 10.09$  mg of vitamin C equivalents/g sample, respectively.

**Initial Quality of VOO:** PV, AV, Free acidity,  $K_{232}$ , and  $K_{268}$  of VOO were  $4.45 \pm 0.68$  meq  $O_2$ /Kg,  $3.62 \pm 0.12$  mg/kg,  $1.69 \pm 0.22\%$ ,  $1.67 \pm 0.25$ , and  $0.16 \pm 0.1$ , respectively, at the beginning of the storage period. Oleic acid was the major fatty acid ( $69.72 \pm 1.70\%$ ) followed by palmitic ( $18.67 \pm 2.68\%$ ), linoleic ( $8.94 \pm 2.20\%$ ), stearic ( $1.72 \pm 0.14\%$ ), and linolenic ( $0.95 \pm 0.63\%$ ) acids. The iodine value of VOO was 83.03. Total phenolic content of VOO was  $290.5 \pm 1.1$  ( $\mu$ g GAE/g).

**Table 1.** Chemical composition of *Zataria multiflora* essential oil using gas chromatography

No.	Compound	Retention index	Relative peak area (%)
1	$\alpha$ -Thujene	924	0.31
2	$\alpha$ -Pinene	932	3.35
3	Camphene	951	0.31
4	$\beta$ -Pinene	975	0.05
5	Myrcene	989	2.46
6	3-Octanol	993	0.07
7	$\alpha$ -Phellandrene	1004	0.34
8	$\alpha$ -Terpinene	1015	2.21
9	<i>p</i> -Cymene	1025	11.74
10	Limonene	1028	0.63
11	$\gamma$ -Terpinene	1058	6.06
12	cis-Sabinene hydrate	1065	0.28
13	Linalool	1098	1.31
14	Borneol	1164	0.95
15	Terpinene-4-ol	1175	1.87
16	$\alpha$ -Terpineol	1189	0.88
17	Thymol	1291	52.8
18	Carvacrol	1305	10.33
19	Thymol acetate	1354	0.92
20	Carvacrol acetate	1371	0.15
21	( <i>E</i> )-Caryophyllene	1417	1.98
22	Aromadendrene	1436	0.29
23	$\alpha$ -Humulene	1451	0.18
24	allo-Aromadendrene	1458	0.25
25	Viridiflorene	1492	0.27

## Oxidative Stability of VOO during Accelerated Storage

**PV, AV, and TV:** Changes in PVs of VOO samples during storage are illustrated in Figure 1a and Figure 1d, respectively. PVs measured in the control were higher than those samples containing *Z. multiflora* EO and BHT during the storage period. The PV of the control increased considerably at the early stages of the storage period, but after 12 days of storage, the PV of the control began to decrease. In fact, after 12 days of storage, the formation of hydroperoxides was slower than their decomposition into secondary oxidation products. In samples containing *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol, it was observed that the PV reached  $12.88 \pm 0.54$ ,  $12.46 \pm 0.06$ , and  $18.10 \pm 0.14$  meq  $O_2$ /kg, respectively, after 16 days storage. The PVs of samples containing antioxidants without CA were significantly higher than those with CA.


AVs of VOO samples are presented in Figure 1b and Figure 1e. The AV of the control sample increased gradually after 8 days of storage. Sharp increase in the AV of the control sample was observed after 12 days of storage. These findings support the idea that hydroperoxides start to decompose and form secondary oxidation products after 12 days of storage (26). BHT, *Z. multiflora* EO, and  $\alpha$ -tocopherol significantly inhibited the formation

of secondary oxidation products in comparison with the control sample. By the end of the storage period, the corresponding percentage of inhibition by *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were 19.87%, 23.99%, and 14.59%, respectively, compared to the control. In the case of samples containing antioxidants with CA, the corresponding percentage of inhibition performed by *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were 22.27%, 28.79%, and 16.51%, respectively, compared to the control.

TVs of VOO samples are illustrated in Figure 1c and Figure 1f. TVs of all samples were increased during storage. *Z. multiflora* EO and BHT were the most stable during the storage period. The effect of *Z. multiflora* EO was comparable to those of BHT when lowering the TV of VOO. The  $\alpha$ -tocopherol was less effective than BHT and *Z. multiflora* EO.

Antioxidant indices of VOO samples are shown in Table 2. *Z. multiflora* EO significantly increased the IP of VOO in comparison with that of the control. No significant difference was observed between IP of samples containing BHT with CA and those containing *Z. multiflora* EO with CA. Therefore, the mixture of *Z. multiflora* EO and CA can be used to replace synthetic antioxidants, i.e. BHT. The  $\alpha$ -tocopherol significantly increased the IPs of VOO in comparison with that of the control, but they were less effective than *Z. multiflora* EO and BHT.

According to PF results, the BHT and *Z. multiflora* EO, in addition to the mixtures of these antioxidants with CA, exhibited the highest PF in VOO during storage period. The  $\alpha$ -tocopherol exhibited a lower PF than the value exhibited by BHT and *Z. multiflora* EO.



**Figure 1.** Effect of *Zataria multiflora* essential oil, BHT, and  $\alpha$ -tocopherol on peroxide values, anisidine values, and TOTOX values of virgin olive oil samples with (d-f) and without (a-c) citric acid.

**Table 2.** Antioxidant indices and synergism activity of virgin olive oil samples during storage period

Treatment	Induction period (day)	Protection factor	Antioxidant activity	Synergism activity (%)
<b>Without citric acid</b>				
Control	7.56±0.17 <sup>d*</sup>	1.00±0.00 <sup>b</sup>	0.00±0.00 <sup>f</sup>	-
BHT	30.84±1.28 <sup>a</sup>	4.07±0.08 <sup>a</sup>	307.99±7.77 <sup>a</sup>	-
<i>Zataria multiflora</i> essential oil	28.21±0.88 <sup>b</sup>	3.73±0.20 <sup>a</sup>	27.31±0.33 <sup>e</sup>	-
$\alpha$ -Tocopherol	16.20±0.14 <sup>c</sup>	2.13±0.04 <sup>b</sup>	113.22±5.06 <sup>c</sup>	-
<b>With citric acid</b>				
Control	8.12±0.03 <sup>d</sup>	1.00±0.00 <sup>b</sup>	0.00±0.00 <sup>f</sup>	-
BHT	32.00±0.99 <sup>a</sup>	3.93±1.06 <sup>a</sup>	146.91±5.23 <sup>b</sup>	2.42±0.72 <sup>ab</sup>
<i>Zataria multiflora</i> essential oil	29.83±0.63 <sup>ab</sup>	3.67±0.06 <sup>a</sup>	24.29±0.54 <sup>e</sup>	4.74±0.69 <sup>a</sup>
$\alpha$ -Tocopherol	16.81±0.24 <sup>c</sup>	2.00±0.01 <sup>b</sup>	53.44±1.02 <sup>d</sup>	1.28±0.71 <sup>b</sup>


\*Values are presented as mean  $\pm$  standard deviation. In each column, means with different letters are significantly different ( $P < 0.05$ ).

Antioxidant activity (AA) of BHT and  $\alpha$ -tocopherol was significantly higher than that of *Z. multiflora* EO. This may be due to the fact that AA depends on the concentration of antioxidant (25). The AAs of samples containing BHT and  $\alpha$ -tocopherol without CA were significantly higher than those with CA, but there was no significant difference between the AA of samples containing *Z. multiflora* EO – samples without CA and those combined with CA.

Synergism activities of tested antioxidants with CA are shown in Table 2. The synergistic activities of CA with *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were 4.74, 2.42, and 1.28%, respectively. Thus, when the tested antioxidants were combined with CA, they showed a synergism in VOO.

**K<sub>232</sub> and K<sub>268</sub>:** K<sub>232</sub> and K<sub>268</sub> values of VOO samples during the storage period are shown in Figure 2. Similar to PV results, K<sub>232</sub> and K<sub>268</sub> values of samples containing *Z. multiflora* EO was lower than samples of the control and samples containing  $\alpha$ -tocopherol. *Z.*

*multiflora* EO delayed the formation of conjugated dienes and trienes to the same extent as BHT did. The CA showed a low synergistic activity with the tested antioxidants in lowering the K<sub>232</sub> and K<sub>268</sub>. The times required to reach the upper permitted limits of K<sub>232</sub> and K<sub>268</sub> for VOO samples during the storage period are shown in Table 3. The durations of time that were required to reach the upper permitted limit of K<sub>232</sub> and IP correlated directly with samples containing CA ( $R^2 = 0.998$ ;  $y = 0.796x + 0.160$ ) and also those without CA ( $R^2 = 0.989$ ;  $y = 0.912x + 0.251$ ). There was a direct correlation between the durations of time that were required to reach the upper permitted limit of K<sub>268</sub> and IP for samples with CA ( $R^2 = 0.926$ ;  $y = 3.651x - 7.304$ ) and those without CA ( $R^2 = 0.997$ ;  $y = 4.597 - 9.983$ ). In addition, the durations of time that were required to reach the upper permitted limit of K<sub>232</sub> and K<sub>268</sub> correlated directly with samples containing CA ( $R^2 = 0.964$ ;  $y = 3.301x - 6.184$ ) and those without CA ( $R^2 = 0.991$ ;  $y = 3.670x - 6.343$ ).



**Figure 2.** Effect of *Zataria multiflora* essential oil, BHT, and  $\alpha$ -tocopherol on K<sub>232</sub> and K<sub>268</sub> values of virgin olive oil samples with (c, d) and without (a, b) citric acid.

**Table 3.** Time required to reach the upper permitted limits of  $K_{232}$  and  $K_{268}$  for virgin olive oil samples during storage period

Treatment	$K_{232}$ upper legal limit (day)	$K_{268}$ upper legal limit (day)	Relative parameters		
			IP/ $K_{232}$	IP/ $K_{268}$	$K_{232}/ K_{268}$
<b>Without citric acid</b>					
Control	7.99±2.11 <sup>d</sup>	3.77±0.90 <sup>d</sup>	0.98±0.24	2.05±0.44	2.11±0.06
BHT	26.49±0.67 <sup>ab</sup>	8.71±0.12 <sup>abc</sup>	1.15±0.01	3.53±0.35	3.06±0.31
<i>Zataria multiflora</i> essential oil	24.12±0.83 <sup>b</sup>	8.46±0.65 <sup>abcd</sup>	1.17±0.07	3.34±0.16	2.86±0.32
$\alpha$ -Tocopherol	14.01±0.01 <sup>c</sup>	5.76±0.52 <sup>cd</sup>	1.15±0.01	2.82±0.23	2.44±0.22
<b>With citric acid</b>					
Control	8.63±1.16 <sup>d</sup>	4.04±0.44 <sup>d</sup>	0.95±0.13	2.02±0.21	2.10±0.09
BHT	29.77±1.74 <sup>a</sup>	10.79±2.28 <sup>a</sup>	1.07±0.03	3.05±0.51	2.76±0.50
<i>Zataria multiflora</i> essential oil	27.73±2.73 <sup>ab</sup>	9.96±0.10 <sup>ab</sup>	1.10±0.08	2.99±0.24	2.95±0.30
$\alpha$ -Tocopherol	14.02±1.05 <sup>c</sup>	6.98±0.83 <sup>bcd</sup>	1.20±0.07	2.42±0.25	2.01±0.09

\*Values are presented as mean  $\pm$  standard deviation. In each column, means with different letters are significantly different ( $P < 0.05$ ).

**Chlorophyll and Carotenoid Contents:** Chlorophyll and carotenoid contents of VOO samples during the storage period are shown in Table 4. Chlorophyll and carotenoid contents of the control group significantly decreased after 16 days storage. In all samples, carotenoid contents decreased more rapidly during the storage period, indicating that carotenoids are more susceptible to degradation than chlorophylls. BHT and *Z. multiflora* EO significantly protected chlorophyll and carotenoid pigments in the VOO against degradation. As a matter of fact,  $\alpha$ -tocopherol was less effective in protecting chlorophyll and carotenoid pigments of VOO than *Z. multiflora* EO and BHT. In terms of chlorophyll and carotenoid contents, there were no significant differences between samples containing CA and those without CA after 16 days storage.

## Discussion

**Analysis of EO:** The main components of *Z. multiflora* EO were oxygenated monoterpenes (thymol and carvacrol) and monoterpene hydrocarbon

(*p*-cymene and  $\gamma$ -terpinene). Thymol was the predominant compound in the present study, which was also considered as the main component of *Z. multiflora* EO by Saei-Dehkordi *et al.* (27). Also, Golmakani and Rezaei reported that *Z. multiflora* EO is predominantly composed of thymol, carvacrol,  $\gamma$ -terpinene, and *p*-cymene (9).

**Antioxidant Activity of EO:** *Z. multiflora* EO exhibited a significant DPPH<sup>o</sup> scavenging activity. The IC<sub>50</sub> value of *Z. multiflora* EO in the present study (0.06 mg/mL) was similar to those reported by Sharififar *et al.* (30) (0.02 mg/mL). Mishra *et al.* reported that among the main components of thyme EO, thymol showed a high free radical scavenging activity with IC<sub>50</sub> value of 0.109  $\mu$ L.mL<sup>-1</sup>, whereas *p*-cymene, showed moderate radical scavenging capacity with IC<sub>50</sub> value of 19.6  $\mu$ L.mL<sup>-1</sup> (28). Also, according to Yanishlieva *et al.*, thymol showed a higher radical scavenging activity than carvacrol. This is due to the greater steric hindrance of the phenolic group in thymol compared to carvacrol (29).

**Table 4.** Chlorophyll and carotenoid contents (mg/kg) of virgin olive oil samples at the beginning and end of storage period

Treatment	Day 0		Day 16		Relative reduction (%)	
	Chlorophyll	Carotenoid	Chlorophyll	Carotenoid	Chlorophyll	Carotenoid
<b>Without citric acid</b>						
Control	4.70±0.07 <sup>a*</sup>	2.05±0.06 <sup>a</sup>	3.02±0.00 <sup>b</sup>	1.02±0.11 <sup>b</sup>	35.58±0.90 <sup>a</sup>	50.18±3.62 <sup>a</sup>
BHT	4.70±0.07 <sup>a</sup>	2.05±0.06 <sup>a</sup>	4.27±0.07 <sup>a</sup>	1.32±0.01 <sup>a</sup>	9.06±0.13 <sup>b</sup>	35.01±2.36 <sup>b</sup>
<i>Zataria multiflora</i> essential oil	4.68±0.06 <sup>a</sup>	2.05±0.06 <sup>a</sup>	4.15±0.02 <sup>a</sup>	1.30±0.01 <sup>a</sup>	11.43±0.86 <sup>b</sup>	35.57±1.98 <sup>b</sup>
$\alpha$ -Tocopherol	4.70±0.07 <sup>a</sup>	2.05±0.06 <sup>a</sup>	3.17±0.20 <sup>b</sup>	1.24±0.02 <sup>a</sup>	32.42±3.28 <sup>a</sup>	39.43±0.12 <sup>b</sup>
<b>With citric acid</b>						
Control	4.70±0.07 <sup>a</sup>	2.05±0.06 <sup>a</sup>	3.15±0.01 <sup>b</sup>	1.09±0.04 <sup>b</sup>	32.96±0.74 <sup>a</sup>	46.96±0.42 <sup>a</sup>
BHT	4.70±0.07 <sup>a</sup>	2.05±0.06 <sup>a</sup>	4.30±0.05 <sup>a</sup>	1.34±0.01 <sup>a</sup>	8.13±0.16 <sup>b</sup>	33.40±0.60 <sup>b</sup>
<i>Zataria multiflora</i> essential oil	4.68±0.06 <sup>a</sup>	2.05±0.06 <sup>a</sup>	4.27±0.07 <sup>a</sup>	1.34±0.00 <sup>a</sup>	8.96±0.18 <sup>b</sup>	34.63±2.60 <sup>b</sup>
$\alpha$ -Tocopherol	4.70±0.07 <sup>a</sup>	2.05±0.06 <sup>a</sup>	3.28±0.23 <sup>b</sup>	1.25±0.01 <sup>a</sup>	30.14±3.80 <sup>a</sup>	39.08±2.44 <sup>b</sup>

\*Values are presented as mean  $\pm$  standard deviation. In each column, means with different letters are significantly different ( $P < 0.05$ ).

*Z. multiflora* EO showed a significant ability to donate electrons to free radicals, converting them into more stable non-reactive species and terminating the free radical chain reactions. It has been reported that *Z. multiflora* EO showed a significant reducing power. *Z. multiflora* EO showed a higher reducing power than *Bunium persicum* EO. The reducing powers of both EOs were lower than TBHQ (31).

**Initial Quality of VOO:** At the beginning of experiment, VOO met the criteria set by the International Olive Council (IOC) for the virgin category regarding the value of free acidity ( $\leq 2.0$ ) (32). Also, PV,  $K_{232}$ , and  $K_{268}$  were below the upper permitted limit established by IOC for VOO ( $PV \leq 20$  meq  $O_2/kg$ ,  $K_{232} \leq 2.6$ , and  $K_{268} \leq 0.25$ ). Therefore, the initial oxidation level of VOO was low. VOO contains a high ratio of monounsaturated/polyunsaturated fatty acids and a low iodine value. This fact is accompanied by a significant amount of total phenols with potential antioxidant capacities, and it results in a high oxidative stability of the VOO. However, presence of polyunsaturated fatty acids can reduce oxidative stability of VOO.

### Oxidative Stability of VOO during Accelerated Storage

**PV, AV, and TV:** Adding the *Z. multiflora* EO and BHT to the samples significantly decreased the PV of samples in comparison with the control during storage. After 16 days storage, the PV did not exceed the upper limit (20 meq  $O_2/kg$ ) permitted by IOC for VOO (32). This indicates that *Z. multiflora* EO is effective in preventing the formation of hydroperoxides. Similarly, Simandi *et al.* reported that 0.6 % thyme extract (*T. vulgaris*) reduced the sunflower oil oxidation to the same extent as 0.1 % BHT did (33). *Z. multiflora* EO was more effective than  $\alpha$ -tocopherol in lowering the PV of VOO. Samotyja and Malecka reported that  $\alpha$ -tocopherol was less effective in sunflower oil stabilization in comparison with rosemary and blackcurrant seed extracts (34).

*Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were effective against formation of secondary oxidation products, mainly in aldehydic structure. Asensio *et al.* showed that extra virgin olive oil samples added with oregano EO had lower anisidine values than the control during storage (13). Also, Vahidyan *et al.* showed that the effect of *Z. multiflora* EO (500 ppm)

in reducing the formation of secondary oxidation products in mayonnaise formulated with linseed oil was similar to that of BHA (1000 ppm) (35).

*Z. multiflora* EO significantly inhibited the TV of VOO in comparison with the control sample. The antioxidant capacity of an EO depends on its chemical composition. Thymol and carvacrol can react with lipid-free radicals and delay the initiation step of oxidation. Also, they can react with peroxy or alkoxy radicals and prevent the propagation step of oxidation. The inhibitory effect of  $\gamma$ -terpinene against lipid oxidation is due to the presence of a strongly activated methylene group which can compete with the activated methylene in the C-11 of linoleic acid, and inhibit linoleic acid oxidation (36).

Combinations of antioxidants with CA showed a synergistic effect on lowering the TV of VOO. Similarly, Hras *et al.* reported that the synergistic activity of rosemary extract combined with CA was 2.61% in the sunflower oil (14).

The effectiveness of different antioxidants commonly corresponds to increasing the IP. This delay is often expressed as PF (14). *Z. multiflora* EO and BHT and their combination with CA showed a "very high" antioxidant activity ( $PF > 3$ ). Similarly, Bandonien *et al.* reported that sage, sweet grass and tansy extracts exhibited very high ( $PF > 3$ ) antioxidant activity (25).  $\alpha$ -Tocopherol both with and without CA showed "medium" antioxidant activity (PF of 2.0-2.5).

Since *Z. multiflora* EO reduced the VOO oxidation at higher concentrations, *Z. multiflora* showed lower AA than that of BHT. It is commonly known that the maximum permitted levels of synthetic antioxidants are established based on various toxicological parameters. However, these parameters and limitations do not need to be applied to naturally occurring compounds. Therefore, *Z. multiflora* EO can be used at higher levels, compared with those of synthetic counterparts, thereby increasing their antioxidant effectiveness.

**$K_{232}$  and  $K_{268}$ :** During the unsaturated fatty acids oxidation, the rearrangement of the double bonds, produce conjugated dienes and conjugated trienes. Conjugated dienes in fats and oils absorb at 232 nm. Similarly, conjugated trienes absorb at 268 nm (26). *Z. multiflora* EO, BHT, and  $\alpha$ -tocopherol were effective in lowering  $K_{232}$  and  $K_{268}$  of VOO. Ayadi *et al.* reported that extra VOOs treated with thyme,



rosemary and basil exhibited a significant low value of  $K_{232}$  and  $K_{270}$  than that of the control during the storage period (37). In this study, the time needed to reach the upper permitted limits of  $K_{268}$  was shorter than those of  $K_{232}$  and PV. Absorbance at 268 nm is due to the oxidation of linolenic acid in VOO (26). Similarly, Gomez-Alonso *et al.* showed that after 21-month storage of VOO at room temperature, linolenic acid reduced 2.5–2.8 times greater than linoleic acid (38). The time required to reach the upper permitted limit of  $K_{232}$  in samples treated with  $\alpha$ -tocopherol were significantly shorter than the durations required by BHT and *Z. multiflora* EO. However, durations of time that were required to reach the upper legal limit of  $K_{268}$  in samples treated with  $\alpha$ -tocopherol were slightly shorter than the durations required by BHT and *Z. multiflora* EO. This observation shows that the activities of different antioxidants differ depending on the oxidizing substrate.

**Chlorophyll and carotenoid contents:** *Z. multiflora* EO can be proposed as natural additives for the preservation of VOO color quality. Also, Asensio *et al.* reported that chlorophyll and carotenoid contents of extra virgin olive oil samples containing oregano EO was higher than that of control during storage (13).

In this study, effect of *Z. multiflora* EO was investigated on VOO oxidation in comparison with the effects of  $\alpha$ -tocopherol and BHT. Also, the synergistic activity of CA was investigated on natural and synthetic antioxidants. Results showed that *Z. multiflora* EO significantly reduced VOO oxidation. This can be attributed to the presence of several antioxidant compounds in *Z. multiflora* EO. The *Z. multiflora* EO reduced VOO oxidation to the same extent as BHT did. The  $\alpha$ -tocopherol was less effective than *Z. multiflora* EO. Also, CA exhibited a synergistic activity with natural and synthetic antioxidants. Generally, *Z. multiflora* EO can be used as natural antioxidants to reduce VOO oxidation and to preserve the color of VOO. Further research will be required to determine the correlation between the antioxidant activity and the chemical composition of *Z. multiflora* EO.

### Acknowledgement

This research project was financially supported by Shiraz University. We would like to thank the Edible Oil Industries Group of Etko Organization for

providing the VOO. We also thank Mohsen Hamedpour-Darabi for editing the English and the research language of the paper.

### Financial disclosure

This research project was financially supported by Shiraz University.

### References

1. Velasco J, Dobarganes C. Oxidative stability of virgin olive oil. *Eur J Lipid Sci Technol* 2002; 104: 661-76.
2. Morales MT, Przybylski R. Olive Oil Oxidation. In: Harwood J, Aparicio R, editors. *Handbook of Olive Oil*. New York: Springer; 2013: 479-522.
3. Basaga H, Tekkaya C, Acikel F. Antioxidative and free radical scavenging properties of rosemary extract. *LWT-Food Sci Tech* 1997; 30: 105-108.
4. Allam, SS, Mohamed, HMA, Thermal stability of some commercial natural and synthetic antioxidants and their mixtures. *J Food Lipids* 2002; 9: 277-293.
5. Sanhueza J, Nieto S, Valenzuela, A, Thermal stability of some commercial synthetic antioxidants. *J Am Oil Chem Soc* 2000; 77: 933-936.
6. Farag R, Badei A, El Baroty G, Palta J, Nobel P, Influence of thyme and clove essential oils on cottonseed oil oxidation *J Am Oil Chem Soc* 1989; 66: 800-804.
7. Yanishlieva NV, Marinova EM. Antioxidative effectiveness of some natural antioxidants in sunflower oil. *Zeitschrift für Lebensmittel-Untersuchung und Forschung* 1996; 203:220-3.
8. Fazeli MR, Amin G, Attari MMA, Ashtiani H, Jamalifar H, Samadi N. Antimicrobial activities of Iranian sumac and avishan-e shirazi (*Zataria multiflora*) against some food-borne bacteria. *Food Cont* 2007; 18: 646-9.
9. Golmakani MT, Rezaei K. Microwave-assisted hydrodistillation of essential oil from *Zataria multiflora* Boiss. *Eur J Lipid Sci Technol* 2008; 110: 448-54.
10. Mastelic J, Jerkovic I, Blažević I, Poljak-Blaži M, Borović S, Ivančić-Baće I, et al. Comparative study on the antioxidant and biological activities of carvacrol, thymol, and eugenol derivatives. *J Agric Food Chem* 2008; 56: 3989-96.
11. Viuda-Martos M, Ruiz Navajas Y, Sánchez Zapata E, Fernández-López J, Pérez-Álvarez JA. Antioxidant activity of essential oils of five spice plants widely used in a Mediterranean diet. *Flav Frag J*. 2010; 25: 13-9.
12. Hashemi MB, Niakousari M, Saharkhiz MJ, Eskandari MH. Stabilization of sunflower oil with *Carum copticum* Benth & Hook essential oil. *J food Sci Tech* 2014; 51: 142-7.
13. Asensio CM, Nepote V, Grosso NR. Chemical Stability of Extra-Virgin Olive Oil Added with Oregano Essential Oil. *J Food Sci* 2011; 76: S445-S50.

14. Hraš AR, Hadolin M, Knez Ž, Bauman D. Comparison of antioxidative and synergistic effects of rosemary extract with  $\alpha$ -tocopherol, ascorbyl palmitate and citric acid in sunflower oil. *Food Chem* 2000; 71: 229-33.
15. Bellik Y, Selles SMA. In vitro synergistic antioxidant activity of honey-Mentha spicata combination. *Journal of Food Measurement and Characterization* 2017; 11: 1-8.
16. Khajehie N, Golmakani MT, Eblaghi M, Eskandari MH. Evaluating the effects of microwave-assisted hydrodistillation on antifungal and radical scavenging activities of *oliveria decumbens* and *chaerophyllum macropodium* essential oils. *J Food Prot* 2017; 80: 783-791.
17. Farahmand M, Golmakani MT, Mesbahi G, Farahnaky A. Investigating the effects of large-scale processing on phytochemicals and antioxidant activity of pomegranate juice. *J Food Process Preserv* 2016; DOI: 10.1111/jfpp.12792.
18. Apak R, Güçlü K, Özyürek M, Çelik SE. Mechanism of antioxidant capacity assays and the CUPRAC (cupric ion reducing antioxidant capacity) assay. *Microchimica Acta* 2008; 160: 413-9.
19. AOCS. Official Methods and Recommended Practices of the American Oil Chemists' Society, 5th ed. Champaign, Illinois: AOCS Press 2000.
20. Golmakani MT, Mendiola JA, Rezaei K, Ibáñez E. Expanded ethanol with CO<sub>2</sub> and pressurized ethyl lactate to obtain fractions enriched in  $\gamma$ -Linolenic Acid from *Arthrospira platensis* (Spirulina). *J Supercrit Fluid* 2012; 62:109-15.
21. Kyriakidis NB, Katsiloulis T. Calculation of iodine value from measurements of fatty acid methyl esters of some oils: comparison with the relevant American oil chemists society method. *J Am Oil Chem Soc* 2000; 77: 1235-8.
22. Casal S, Malheiro R, Sendas A, Oliveira BP, Pereira JA. Olive oil stability under deep-frying conditions. *Food Chem Toxicol* 2010; 48: 2972-9.
23. Minguez-Mosquera MI, Rejano-Navarro L, Gandul-Rojas B, SanchezGomez AH, Garrido-Fernandez J. Color-pigment correlation in virgin olive oil. *J Am Oil Chem Soc* 1991; 68: 332-6.
24. Bandonien D, Pukalskas A, Venskutonis P, Gruzdien D. Preliminary screening of antioxidant activity of some plant extracts in rapeseed oil. *Food Res Int* 2000; 33:785-91.
25. Antolovich M, Prenzler PD, Patsalides E, McDonald S, Robards K. Methods for testing antioxidant activity. *Analyst* 2002; 127: 183-98.
26. Frankel EN, *Lipid Oxidation*, 2nd ed. Cambridge: Woodhead Publishing Limited 2005.
27. Saei-Dehkordi SS, Tajik H, Moradi M, Khalighi-Sigaroodi F. Chemical composition of essential oils in *Zataria multiflora* Boiss. from different parts of Iran and their radical scavenging and antimicrobial activity. *Food Chem Toxicol* 2010; 48: 1562-7.
28. Mishra PK, Singh P, Prakash B, Kedia A, Dubey NK, Chanotiya C, Assessing essential oil components as plant-based preservatives against fungi that deteriorate herbal raw materials. *Internat Biodet Biodeg* 2013; 80: 16-21.
29. Yanishlieva NV, Marinova EM, Gordon MH, Raneva VG. Antioxidant activity and mechanism of action of thymol and carvacrol in two lipid systems. *Food Chem* 1999; 64: 59-66.
30. Sharififar F, Moshafi M, Mansouri S, Khodashenas M, Khoshnoodi M, In vitro evaluation of antibacterial and antioxidant activities of the essential oil and methanol extract of endemic *Zataria multiflora* Boiss. *Food Control* 2007; 18: 800-805.
31. Zangiabadi M, Sahari M, Barzegar M, Naghdi Badi H, *Zataria multiflora* and *Bunium persicum* essential oils as two natural antioxidants. *J Med Plants* 2012; 1: 8-21.
32. IOC. Trade Standard Applying to Olive Oils and Olive-pomace oils. Decision COI/T.15/NC No 3/Rev. 8. Madrid, Spain: International Olive Council 2015.
33. Simandi B, Hajdu V, Peredi K, Czukor B, Nobik-Kovacs A, Kery A. Antioxidant activity of pilot-plant alcoholic and supercritical carbon dioxide extracts of thyme. *Eur J Lipid Sci Technol* 2001; 103:355-8.
34. Samotyja U, Małecka M. Antioxidant activity of blackcurrant seeds extract and rosemary extracts in soybean oil. *Eur J Lipid Sci Technol* 2010; 112:1331-6.
35. Vahidyan H, Sahari M, Barzegar M, Naghdi Badi H, Application of *Zataria multiflora* Boiss. and *Satureja hortensis* L. essential oils as two natural antioxidants in mayonnaise formulated with linseed oil. *J Med Plants* 2012; 3:69-79.
36. Ruberto G, Baratta MT. Antioxidant activity of selected essential oil components in two lipid model systems. *Food Chem* 2000; 69: 167-74.
37. Ayadi M, Grati-Kamoun N, Attia H. Physico-chemical change and heat stability of extra virgin olive oils flavoured by selected Tunisian aromatic plants. *Food Chem Toxicol* 2009; 47:2613-9.
38. Gómez-Alonso S, Mancebo-Campos V, Salvador MD, Fregapane G. Evolution of major and minor components and oxidation indices of virgin olive oil during 21 months storage at room temperature. *Food Chem* 2007; 100:36-42.