

Review Article

Diet Quality and Its Assessment Methods: A Narrative Review

Sayed Reza Hojati¹, Fatemeh Rostamian¹, Sareh Eghtesad², Mojtaba Farjam³, Jalaledin Mirzay Razzaz^{4,*}, Seyed Hossein Davoodi⁵, Elham Ehrampoush³, Reza Homayounfar^{6,*}

- 1- Student Research Committee, Fasa University of Medical Sciences, Fasa, Iran
- 2- Liver and Pancreatobiliary Diseases Research Center, Digestive Diseases Research Institute, Tehran University of Medical Sciences, Tehran, Iran
- 3- Noncommunicable Diseases Research Center, Fasa University of Medical Sciences, Fasa, Iran
- 4- Department of Community Nutrition, Faculty of Nutrition and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, Iran
- 5- Department of Clinical Nutrition and Dietetics, National Nutrition and Food Technology Research Institute, Faculty of Nutrition Science and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran
- 6- National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Received: May 2024 Accepted: June 2024

ABSTRACT

Background and Objectives: Most studies in the field of nutrition focus on diet quantity and its relationships with health and diseases. However, effects of diet quality on disease development have not been investigated. Diet quality is a novel concept but its assessment is not carried out frequently because of difficulties in carrying out accurate assessments. Several indices have been developed to assess the overall diet quality of individuals and populations. Indices are diverse and each is used in a specific condition to achieve a certain goal. This article aimed to introduce the most importantly useful diet-quality indices and their uses to encourage their widespread uses.

Materials and Methods: Using PubMed, Scopus, Elsevier and Google Scholar databases, keywords such as "diet", "diet quality", "diet quality index", "diet quality score", "nutritional assessment", "food intake pattern" and "nutritional pattern" were used to gather data from 45 associated articles, all of which were used in this manuscript to introduce in detail and assess 13 diet quality indices.

Results: For each index, previous research, current uses and use criteria were reported.

Conclusions: To assess risks of chronic diseases and mortality from cardiovascular diseases, diet quality index, healthy eating index, alternative healthy eating index and healthy diet indicator can be used. For predicting breast cancer risks, diet quality index-revised and alternative healthy eating index can be benefitted. Assessing diet quality parallel to diet quantity renders a further complete assessment of the relationships between diet and disease.

Keywords: Nutritional assessment, Diet quality index, Diet quality, Diet quality score

Highlights

- Assessing diet quality parallel to diet quantity renders a further assessment of the relationships between diet and disease.
- For assessing the risk and mortality from CVDs, AHEI is more reliable because
- To predict mortality from food patterns, MDS versions are the best choices.

Introduction

When the term "diet" is encountered, most people think of a routine diet program for losing weight or disease control and prevention. In contrast, "diet" is a general term and in fact means the total quantity of food and drink habitually consumed by an individual or population. Diet is one of the most important aspects of individuals' lives. Diet and its associated nutrients highly affect longevity (1-4), weight (5-9), development of diseases (10-12) and disorders such as depression (13-16). A healthy diet contains adequate quantities of all macronutrients (proteins, carbohydrates and beneficial fats), all of which are necessary for the human health. In contrast, an unhealthy diet can result in poor growth (17), decreased fertility (18, 19), decreased immunity and increased risk of diseases (20). Comparing and differentiating healthy diets from unhealthy diets is part of a subject called "diet quality".

Recently, diet quality has attracted attention of researchers in the field of nutrition (21). The term "diet quality" is widespread and complex. This term assesses personal food tastes and food diversity of various cultures (22). Diet quality encompasses the diversity, balance and healthfulness of a diet. A diet is addressed to include good quality when it provides the adequate quantities of energy and nutrients needed for growth as well as having a healthy and active life. Food diversity is critical in meeting people's nutritional requirements. It is important to pay attention to diet quality to maintain physiological homeostasis, enhance development and physical activity and prevent the spread of infectious diseases (23).

analytical methods are available to Several quantitatively assess diet quality, such as assessing dietary recall questionnaires for a specific time and assessing food intake quantities. It is difficult to assess diet quality; however, as it includes a detailed assessment of the types of foods and number and size of the meals and their frequency. Diet quality may be associated to eating styles, and other food habits. In addition, snacking microbiological quality and ingredient details of a single food such as meat, dairy or vegetables may be variable and must be addressed (22). A number of assessment indices are currently available, the use of which is increasing. These indices are used to assess epidemiological relationships between diets and their associated health outcomes (24). However, there is no agreements on how to define diet quality or how to set a specific framework for indices for its assessment (21). Therefore, the aim of this study was to compare available diet quality-assessment indices from the literature and assess their weaknesses and strengths to present the most useful indices in assessing diet quality.

Materials and Methods

Through the PubMed, Scopus, Elsevier and Google Scholar databases, keywords such as "diet", "diet quality", "diet quality index", "diet quality score", "nutritional assessment", "food intake pattern" and "nutritional pattern" were used to gather data of 45 associated articles, all of which were used in this manuscript to introduce in detail and assess 13 diet quality indices.

Results

Indices of diet quality assessment

The 13 diet-quality indices assessed in this study as well as their use are listed in Table 1 and further discussed in the following paragraphs.

Diet quality index

Diet quality index (DQI) was first published in 1994 by Patterson (25). This index is not applicable to diseases that are created or treated with a single food or nutrient, but is used to assess the risk of chronic diseases associated with food patterns (25). It is used to investigate mortality from cardiovascular diseases (CVDs), but not other diseases such as cancers. This index consists of eight components that include six micro/macronutrients and two food groups. The micro/macronutrients include total fat, proteins, saturated fatty acids (SFAs), cholesterol, calcium and sodium. Food groups included fruits and vegetables (combined) and the complex carbohydrate groups. The components were scored as follows: if the quantity of a component was equal to, close to and far from the suggested value, it received a score of 0, 1 and 2, respectively. The possible score range was 0–16 (24). The lower the score, the better the diet quality and vice versa (Table 2).

Diet Quality Index-Revised

The original version of DQI was revised in 1999 and diet quality index-revised (DQI-R) was introduced (28). This new index was updated to better reflect the diet quality and improve diet regulation. The assessment of dietary balance and diversity was then added to this index (29). The DQI-R consists of ten components, including cholesterol, fatty acids, total fat, vegetables, fruits, cereals, iron, calcium, dietary diversity score (DDS) and balanced diet score. The first three components represent macronutrient intake. The next three components represent use of the highlighted groups. Calcium and iron components are assessed based on dietary reference intake (DRI). The last two components represent importance of food diversity and controlled consumption of salt, fats and alcoholic beverages and balanced consumption of sugar. The score assigned to each component ranged 0-10, with an overall score of 0–100. In contrast to the original DQI, the higher the score of DQI-R, the better the diet quality (28). It is believed that the 100-point scale made the interpretation of DQI-R easier (29). Differences with the

Nutrition and Food Sciences Research 42 Vol 11, No 4, Oct-Dec 2024

original DQI are as follows: In the DQI, vegetables and fruits were combined into one group, but are separated in the DQI-R. Iron intake has been added to this index, while protein has been eliminated. Food balance and diversity have been added to the DQI-R (24). The novel index

revealed the relationship between this index and plasma biomarkers that reflect nutrient intake (30). Fung used this index to predict breast cancer risks (31). In another study, Fung showed that the DQI-R did not correlate with plasma CVD biomarkers (Table 3) (32).

Table 1. An overall view of the indices

	Index	Components	Details	References
1	DQI (Diet Quality Index)	Total fat, Cholesterol, Saturated fatty acid, Vegetables and fruit and, Protein, Complex carbohydrates, Sodium, Calcium	assesses the risk of chronic diseases associated with the food pattern, investigates mortality from cardiovascular diseases	(24-27, 50)
2	DQI-R (Diet Quality Index- Revised)	total fat, fatty acids, cholesterol, fruits, vegetables, cereals, calcium, iron, dietary diversity score (DDS) and balanced diet score	The relationship between this index and plasma biomarkers was studied, which indicate nutrient intake. predict breast cancer risk	(24, 27-30, 32, 51)
3	DQI-I (Diet Quality Index- International)	diversity, adequacy, moderation and balance	compares diet quality among nations, compares the diet quality in different cultures and to carry out a global monitoring and investigation on the diet quality	(33)
4	HEI (Healthy Eating Index)	fruits, vegetables, grains, milk, meat and beans, Total fat, Saturated fatty acid, Cholesterol, Sodium, Variety	shows changes in food patterns and also promotes health status and proper nutrition education	(28, 34)
5	Alternative Healthy Eating Index (AHEI)	vegetables, fruits, nuts and soybeans, white-red meat ratio, the percentage of energy derived from trans-fat (trans-unsaturated fatty acids), cereal fiber, polyunsaturated fatty acids (PUFA)-saturated fat ratio, daily alcohol consumption and the duration of taking multivitamins	confirms the benefits of unsaturated fats, distinguishes quality in different food groups	(27, 36)
6	HDI (Healthy Diet Indicator)	Protein, Complex carbohydrates, Dietary fiber, Saturated fatty acids, Polyunsaturated fatty acid, Pulses/ nuts/seeds, Fruits and vegetables, Mono-and disaccharides, Cholesterol	It was used for the prevention of chronic diseases, It was associated with a reduction in all mortality factors	(37, 38)
7	Healthy Food Index (HFI)	lack of taking margarine / butter / fats, consuming raw or boiled vegetables at least for one time, using at least one white bread or large rye bread and consuming fruits at least for one time	Investigates the relationship between diet quality and all mortality factors	(19, 33, 34)
8	Mediterranean Diet Score (MDS)	high MUFA to saturated fats proportion, high consumption of legumes, high intake of vegetables, fruits, grains, moderate consumption of alcohol, low intake of milk and dairy products and low intake of meat and meat products	Evaluates the overall diet quality according to the traditional Mediterranean diet in Mediterranean populations	(24, 26, 27, 41)
9	Mediterranean Adequacy Index (MAI)	Mediterranean food group includes cereals, legumes, vegetables, fruits, starches, fish, MUFA and liquor. The non-Mediterranean food group includes milk/dairy products, meat/poultry, eggs, sugar and saturated fats	Total Mediterranean food groups were divided by the total non- Mediterranean food groups. It related to mortality rates	(24, 44)
10	Food Variety Score (FVS)	the number of different foods that a person consumes in a given time period	consumption frequency and consumption rate are not investigated	(46, 47)
11	Dietary Diversity Score (DDS)	the number of food groups used usually based on food pyramid groups	widely used by researchers in the nutrition field	(47, 48)
12	Nutritional Adequacy Ratio (NAR)	indicates the adequacy of nutrients intake based on the recommended dietary allowance (RDA)	the amount of nutrient intake is divided by the recommended amount and is reported as a percentage	(26, 45, 49)
13	Mean Adequacy Ratio (MAR)	By calculating of the average NARs	presented as a general index for nutrient adequacy	(26, 45, 49)

Table 2. Component guide table and scoring method of diet quality index (26, 27)

		<30 energy %	0
1	Total fat	30-40 energy %	1
		>40 energy %	2
		<10 energy%	0
2	Saturated fatty acid	10-13energy %	1
		>13energy %	2
		<300 mg	0
3	Cholesterol	300-400 mg	1
		>400 mg	2
		5+ serving	0
4	Fruit and vegetables	3-4 serving	1
		0-2 serving	2
	Complex carbohydrates	6+ serving	0
5		4-5 serving	1
		0-3 serving	2
		≤100% RDA	0
6	Protein	100-150% RDA	1
		≥150% RDA	2
		<2400 mg	0
7	Sodium	2400-3400 mg	1
		>3400 mg	2
		≥RDA	0
8	Calcium	2/3 RDA	1
		<2/3 RDA	2

Table 3. Component guide table and scoring method of diet quality index-revised (27, 28)

		≤ 30 energy %	10
1	Total fat $\leq 30\%$	>30 energy %	5
		> 40 energy %	0
	Saturated fatty acids	≤ 10 energy %	10
2	≤ 10% energy intake	10≤13 energy %	5
	≥ 10% energy intake	>13 energy %	0
		≤300 mg	10
3	Cholesterol	>300, ≤400 mg	5
		>400 mg	0
		≥ 100%	
4	2-4 serving fruit per day	99%-50%	$0-10^{*}$
	• • •	< 50%	
		≥ 100%	
5	3-5 serving vegetables per day	99%-50%	$0-10^{*}$
		<50%	
		≥ 100%	
6	6-11 serving grains per day	99%-50%	$0-10^{*}$
	4 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 -	<50%	
		≥ 100%	
7	Calcium intake	99%-50%	$0-10^{*}$
	Cultum munic	<50%	
		≥ 100%	
8	Iron intake	99%-50%	$0-10^{*}$
	non make	<50%	
		≥6	
9	Dietary diversity score	 ≥3 ,<6	0-10
	Dietary diversity score	<3	v - v
		 ≥7	
10	Dietary moderation score		0-10
	Diemij modelmion score	, · · · · <4	0 10
*Deper	nding on energy intake	**	
	<u> </u>		

Nutrition and Food Sciences Research 44 Vol 11, No 4, Oct-Dec 2024

Diet quality index-international

Diet quality index-international (DQI-I) was proposed to fill for the lack of an index to compare diet quality among nations. This index was released in 2003 to compare diet quality in various cultures and to carry out global monitoring and investigation on diet quality. The

DQI-I has outperformed previous versions of DQI and the healthy eating index (HEI) thus appearing to be more reliable. It includes many aspects of diet quality, measuring diversity, adequacy, moderation and balance (Table 4) (33).

Table 4. Component guide table and scoring method of diet quality index-international (33)

A	Variation		0-20
1	All food groups (fish/poultry/meat/eggs; beans/dairy;	\geq 1 serving from each food group/d = 15	0-15
	grain; vegetable; fruit)	Any 1 food group missed/ $d = 12$	
		Any 2 food groups missed/ $d = 9$	
		Any 3 food groups missed/ $d = 6$	
		\geq 4 food groups missed/d = 3	
		None of any food groups $= 0$	
2	Within-group diversity for protein source (fish,	\geq 3 different origins/d = 5	0-5
	poultry, meat, eggs, dairy, beans)	2 different origins $/d = 3$	
		From 1 origins $d = 1$	
		None = 0	
3	Adequacy		0-40
3	Vegetable group	\geq 3–5 servings/d = 5	0-5
		0 servings/d = 0	
4	Fruit group	\geq 2–4 servings/d = 5	0-5
		0 servings/d = 0	
5	Grain group	\geq 6–11 servings/d = 5	0-5
		0 servings/d = 0	
6	Fiber	$\geq 20-30 \text{ g/d} = 5$	0-5
		0 g/d = 0	
7	Protein	$\geq 10\%$ of energy/d = 5	0-5
		0% of energy/d = 0	
8	Iron	$\geq 100\%$ of RDA (AI)/d = 5	0-5
		0% of RDA (AI)/d = 0	
9	Calcium	≥100% AI/d = 5	0-5
		0% AI/d = 0	
10	Vitamin C	$\geq 100\%$ of RDA (RNI)/d = 5	0-5
		0% of RDA (RNI)/d = 0	
C	Restraint		0-30
11	Total fat	≤20% total energy/d = 6	0-6
		>20-30% total energy/d = 3	
		>30% total energy/d = 0	
12	Saturated fat	\leq 7% total energy/d = 6	0-6
		>7-10% total energy/d = 3	
		>10% total energy/d = 0	
13	Cholesterol	$\leq 300 \text{ mg/d} = 6$	0-6
		>300-400 mg/d = 3	
		>400 mg/d = 0	
14	Sodium	$\leq 2400 \text{ mg/d} = 6$	0-6
		$\frac{1}{2400-3400}$ mg/d = 3	
		>3400 mg/d = 0	
15	Empty calorie foods	\leq 3% of total energy/d = 6	0-6
	* *	>3-10% of total energy/d = 3	
		>10% of total energy/d = 0	
D	Total balance		0-10
16	Macro-nutrient ratio (carbohydrate:protein:fat)	55 ~ 65:10 ~ 15:15 ~ 25 = 6	0-6
	. , ,	52 ~ 68:9 ~ 16:13 ~ 27 = 4	
		$50 \sim 70.8 \sim 17.12 \sim 30 = 2$	
		Otherwise = 0	
17	Fatty acid ratio (PUFA:MUFA:SFA)	$P/S = 1 \sim 1.5$ and $M/S = 1 \sim 1.5 = 4$	0-4
	,	Else if $P/S = 0.8 \sim 1.7$ and $M/S = 0.8 \sim 1.7 = 2$.

Vol 11, No 4, Oct-Dec 2024

Total score of these four components is summed, with a maximum score of 100. The DQI-I includes its own grouping. For example, empty-calorie foods, which have not been addressed in other indices, are a food group in this index. The DQI-I requires a lot of diet information, which is collected through several-day diet recalls, making it time consuming to carry out (33).

Healthy eating index

The HEI, introduced by Kennedy (34), is an assay that briefly assesses diet quality and can be used to show changes in food patterns and promote health status and appropriate nutrition education. The US Department of Agriculture (USDA) includes a plan to use the HEI as a base for nutrition promotion programs (34). Components of this index are divided into two categories of adequacy and moderation. Adequacy subcategories are addressed to ensure adequate intake of nutrients, while the intake of nutrients should be limited in moderation subcategories. Technically, HEI consists of ten components of determining compliance with the number of servings recommended by the Food Guide Pyramid regarding five major food groups (grains, vegetables, fruits, milks, meats and beans); the total intake of sodium; saturated fats and total intake of fat as part of the energy intake; quantity of cholesterol intake; and diet diversity. Each component is assigned a score of 0–10 with the final score of 0–100, indicating the worst and the best diets, respectively (Table 5) (28).

Alternative healthy eating index

Alternative healthy eating index (AHEI) was assessed for disease risk prediction, compared to HEI. The AHEI verifies benefits of unsaturated fats (USFs) (similar to HEI), yet distinguishes quality in various food groups and removes potatoes and its products from the vegetables group. It consists of nine components, including vegetables, fruits, nuts and soybeans, white-red meat ratios, energy proportions derived from trans-fat [transunsaturated fatty acids (trans-UFAs)], cereal fibers, polyunsaturated fatty acids (PUFAs)-saturated fat (SF) ratios, daily alcohol consumption and duration of multivitamin use. The total scores of all components show the AHEI score, ranging from 87.5 (the best diet) to 2.5 (the worst diet) (24). Results of a study showed that the incidence of breast cancer decreased by 11% with a 10% increase in AHEI (31, 35). Another study showed that AHEI predicts risks of chronic diseases better than those the HEI does, which is due to the strong inverse relationships between AHEI and CVDs (Table 6) (36).

Table 5. Component guide table and scoring method of healthy eating index (26–28, 34)

	Adequacy		•
1	Grain	6-11 serving	0-10
2	Vegetables	3-5 serving	0-10
3	Fruits	2-4 serving	0-10
4	Milk	2-3 serving	0-10
5	Meat and beans	2-3 serving	0-10
	Moderation		
6	Total fat	<30 % energy	0-10
7	Saturated fatty acid	<10 % energy	0-10
8	Cholesterol	<300 mg	0-10
9	Sodium	<2400 mg	0-10
10	Variety	16 different food items/3d	0-10

The criteria depend on energy intake. 0 servings score=0

Table 6. Component guide table and scoring method of alternative healthy eating index (27, 36)

1	Vegetables	0 serving	0
	Vegetables	5 serving	10
2	Fruits	0 serving	0
	Truits	4 serving	10
3	Nuts and soy protein	0 serving	0
3	Nuts and soy protein	1 serving	10
1	Ratio of white to red meat	0	0
4	Ratio of white to fed meat	4	10
	C1 £1/J	0	0
3	Cereal fiber g/d	15	10
	Tr. C.	≥4 energy %	0
6	Trans fat	≤0.5 energy%	10
7	Delympostymated, activated fatty and matic	≤0.1	0
/	Polyunsaturated: saturated fatty acid ratio	≥1	10
8	Duration of multivitamins use	< 5 years	0
0	Duration of multivitainins use	≥ 5 years	10
		Men: 0 or >3.5	0
	Alcohol	Women: $0 \text{ or } > 2.5$	U
9		Men: 1.5-2.5	10
		Women: 0.5-1.5	10

Nutrition and Food Sciences Research Vol 11, No 4, Oct-Dec 2024

Healthy diet indicator

Healthy diet indicator (HDI) is based on the dietary recommendations of World Health Organization (WHO) for chronic disease prevention. This index is associated with a decrease in all mortality factors and, in particular, significantly decreased mortality from CVDs (37). It has been shown that the rate of cognitive impairment decreases with increases in HDI (38). This index consists of nine components. The scoring method is as follows: In each component "Score 1" is assigned for intake rates that are within the suggested range and "Score 0" for anything otherwise, adding up to a possible score of 9. The HDI is most commonly used to assess the relationship between mortality and food patterns (Table 7) (37).

Healthy food index

Healthy food index (HFI) was based on previous diet quality assessment indices in 2001. It consists of four components; to which, a score of 1 is assigned in case a desired behavior is carried out daily, otherwise a score of 0 is received. Components include lack of margarine/butter/fat use, consuming raw or boiled vegetables at least once, using at least one white bread or large rye bread and consuming fruits at least once. The total scores range 0-4 and a higher score indicates a better diet quality. This index has been used to investigate the relationship between diet quality and all mortality factors in various research and shown to include an inverse relationship (Table 8) (24, 39, 40).

Mediterranean diet score

Mediterranean diet score (MDS) assesses overall diet quality based on the traditional Mediterranean diet in Mediterranean populations. It was first introduced by Trichopoulou in 1995 (41). Later, new versions were released over the years. Although there are slight differences between the various versions, they include significant effects on scoring, classification and relationship with health outcomes. Further studies are needed to assess which version of MDS is the best choice for investigating the relationship between diet quality and health outcomes (26).

MDS (1): The basic or original version of MDS consists of eight components, including high MUFA to SF proportions, high intakes of legumes, grains, vegetables and fruits, low intakes of milk and dairy products and moderate intakes of alcohol, meat and meat products. Scoring is carried out based on the average intake for each gender. If intake of the first five components exceeds the median, a score of 1 is assigned to that component and if it is less than the average, a score of 0 is assigned. For the rest of three components, scores 1 and 0 are assigned if the quantity consumed is less and greater than the average value, respectively. The possible score range is 0-8, with higher scores indicating better quality. This index was used to investigate the relationship between diet and survival. There are reports that 1 score increase in MDS decreases the overall mortality in elderly people by 17% (Table 9) (24, 26, 27, 41).

Table 7. Component guide table and scoring method of healthy diet indicator (26, 27, 37)

1	Saturated fatty acids	0-10 energy %	1
2	Polyunsaturated fatty acid	3-7 energy %	1
3	protein	10-15 % energy	1
4	Complex carbohydrates	50-70 % energy	1
5	Dietary fiber (g)	27-40 % energy	1
6	Fruits and vegetables(g)	>400 g/d	1
7	Pulses, nuts, seeds (g)	>30 g/d	1
8	Mono-and disaccharides	0-10 energy %	1
9	Cholesterol(mg)	0-300 mg/d	1

If percent or quantities are not in the ranges=0

Table 8. Component guide table and scoring method of healthy food index

1	Not consuming margarine, butter or lard	If met daily	1
		If not met daily	0
2	Consumption of boiled or raw vegetables at least once	If met daily	1
		If not met daily	0
3	Consumption of coarse rye or white bread at least once	If met daily	1
		If not met daily	0
4	Consumption of fruit at least once	If met daily	1
		If not met daily	0

Vol 11, No 4, Oct-Dec 2024

47

Nutrition and Food Sciences Research

Table 9. Component guide table and scoring method of all Mediterranean diet score versions

	MDS (1)	MDS (2)	MDS (3)	MDS (4)	Criteria	a
1	Monounsaturated: saturated	Monounsaturated: saturated	Monounsaturated: saturated	Monounsaturated: saturated	<median< td=""><td>0</td></median<>	0
1	fatty acids ratio	fatty acids ratio	fatty acids ratio	fatty acids ratio	≥median	1
2	Legumes	Legumes	_	Legumes, Nuts and Seeds	<median< td=""><td>0</td></median<>	0
	Legumes	Legumes		Degames, Ivats and Beeds	≥median	1
3	Cereals	Cereals	Cereals (starchy roots	Cereals	<median< td=""><td>0</td></median<>	0
	Coronis	Coronis	excluded)	Corouis	≥median	1_
4	Fruits and nuts	Fruits and nuts	Fruits and nuts	Fruits	<median< td=""><td>0</td></median<>	0
	Traits and nats	Traits and nats	Traits and nats	Truits	≥median	1
5	Vegetables	Vegetables	Vegetables (starchy roots	Vegetables and Potatoes	<median< td=""><td>0</td></median<>	0
	v egetables	v egetables	included)	v egetables and I otatoes	≥median	1_
6	Meat and meat products	Meat and meat products	Meat and meat products	Meat and poultry	≥median	0
	wicat and meat products	Weat and meat products	Weat and meat products	Meat and pountry	<median< td=""><td>1</td></median<>	1
7	Milk and dairy products Milk and dairy pro	Milk and dairy products	Milk and dairy products	Milk and dairy products	≥median	0
	wink and dairy products	with and daily products	wink and dairy products	wink and dairy products	<median< td=""><td>1</td></median<>	1
8	_	Fish	_	Fish	<median< td=""><td>0</td></median<>	0
		Tish		1 1511	≥median	1
9	Alcohol	_	Alcohol	Alcohol	≥median	0
	Alcohor		Aiconoi	7 Heorioi	<median< td=""><td>1</td></median<>	1
				Men ((10-50 g/d)	1
10	-	Alcohol	-	Women	(5-25 g/d)	1
					otherwise	0

MDS (2): The second version of MDS was released in 2003. It is quite similar to the original version and only a few minor improvements were created. Fish consumption was added as a new component. The scoring system is the same as the original version, with fish consumption receiving a score similar to the first five components. Another change was created for scoring alcohol consumption. If daily alcohol intakes for men and women were 10-50 and 5-25 g, respectively, score 1 is assigned, otherwise score 0 is assigned. The possible score range is 0–9, with score of 9 representing the highest compliance with the Mediterranean diet. This version has been used to investigate the relationship between diet quality and overall mortality rate, A 2-point increase in MDS (2) was associated with a 25% decrease in the overall mortality rate in elderly people (24, 42).

MDS (3): This version of MDS is similar to the first version overall except elimination of legumes, leaving a score range of 0–7. Higher scores indicate better food behaviors. This version is used to investigate the relationship between all types of diet-related deaths. It was shown that the overall mortality rate in the elderly people decreased by 21% with 1-score increase in this version of MDS. Higher scores corresponded with higher serum carotene levels and an inverse relationship between serum carotene level and mortality rate was observed (24, 43).

MDS (4): The fourth version of the MDS differs from the original version in that the legume group was changed to the legumes/nuts/seeds group, the vegetable group was changed to the vegetables/starch groups and the meat and meat products group was changed to the poultry and meat group. Fish was added as a separate group. Nine food groups were created with a total score of 0–9. Results of studies show a clear inverse relationship between this

index or HDI and the Mediterranean adequacy index (MAI) and all mortality factors. In addition, a direct relationship was observed between the consumption of SFAs and mortality (24, 44).

Mediterranean adequacy index

The MAI consists of two versions of one original and another one modified. In the modified version, the total Mediterranean food groups (cereals, legumes, vegetables, fruits, starches, fish, MUFAs and liquors) were divided by the total non-Mediterranean food groups (milk/dairy products, poultry/meat, sugar, eggs and SFs). Vegetable oils are used as an alternative of MUFAs and animal fats and margarines shifted to SFAs in the original version. The food groups are articulate as a total daily energy intake percentage in the original version, while consumption values are based on the daily intake of men and women in the modified version (24, 44). Recent researches have shown that the MAI score to be inversely associated to mortality rates (44).

$$MAI = \frac{Sum \ of \ mediterranean \ food \ groups}{Sum \ of \ non-mediterranean \ food \ groups}$$

Food variety score

Food variety score (FVS) simply is the number of various foods that a person consumes within a specific time. Consumption frequency and consumption rate are not investigated (45). Various studies have used this index (46, 47).

Dietary diversity score

The DDS is based on the number of consumed food groups from the food pyramid groups. For each group, scores 1 and 0 are assigned for consuming and not consuming a food groups, respectively, and the sum of

Nutrition and Food Sciences Research 40 Vol 11, No 4, Oct-Dec 2024

scores is expressed as DDS (45). The FVS and DDS can be addressed equivalent to food group indicators (FGI) (22). The DDS has widely been used by researchers in the nutrition field (47, 48).

Nutritional adequacy ratio

Madden and Yoder published nutritional adequacy ratio (NAR) during a nutrient intake survey in 1972 (49). This ratio indicates the adequacy of nutrient intake based on the dietary reference intakes (DRI), recommended dietary allowances (RDA) or recommended nutrient intakes (RNI) for each age and sex group. In other words, quantity of nutrient intake is divided by the recommended quantity and reported as a percentage (26). The NAR of the energy consumption can be calculated based on the average daily energy requirement or total energy expenditure (TEE). In addition, NAR can be calculated on the basis of reference values for energy derived from various dietary components such as energy from carbohydrates, fats and proteins (45).

Mean adequacy ratio

In addition to NAR, the mean adequacy ratio (MAR) was present as a general index for nutrient adequacy. The MAR is achieved by calculating the average of NARs. To calculate MAR, NARs that are more than 1 or 100% should be truncated at 1 or 100%; thus, high-NAR nutrients do not compensate theoretically low-NAR nutrients. Up-to-date, NAR and MAR indices have been used to investigate diet quality in various studies (26, 45, 49).

MAR (Mean Adequacy Ratio) $= \frac{\sum NAR (each one should be truncated at 1)}{Number of nutrients}$

Discussion

Assessing the diet quality can play a significant role in nutrition-related disease studies, because diseases are not only relevant to nutrient quantity, but they are relevant to the diet quality. Various indices have been formulated to assess diet quality of various populations. Assessing diet quality can play a significant role in nutrition-related diseases to assess the risk of cancers, chronic diseases and all mortality factors. There are specific indices that can be used. In addition, region-specific indices are preferred over general indices to assess diet quality of certain populations.

Conclusion

For assessing the risk of chronic diseases and mortality from CVDs, various indices can be used, but AHEI can be further reliable because it concludes more risk factors than those others do. In the next level, it is better to benefit from DQI and HDI because they show their power in recent studies. To predict breast cancer risk, DQI-R and AHEI can be used but AHEI is used further. To predict mortality from food patterns, MDS versions are the best choices. As studies have revealed that Mediterranean diet is the best diet for people with CVDs, it is better to increase this index in such people. Between MDS versions, MDS (4) is further complete since it includes more diet dimensions than those the others do. The HEI is further useful to educate population how to have appropriate nutrition. It is a good idea to compare diet quality in nations and assess why the prevalence of some diseases is higher in certain countries. Therefore, DQI-I should be used (Table 10).

Table 10. Guide to select the most appropriate index

What are you looking for?	Indexes that can be used
Assessing the risk of chronic diseases associated with the food pattern	DQI, HEI, AHEI, HDI
Investigating mortality from cardiovascular diseases	DQI, HEI, AHEI, HDI
Predicting breast cancer risk	DQI-R, AHEI
Diet quality among nations	DQI-I
Promoting health status and proper nutrition education	HEI
Predicting mortality from food patterns	HDI, HFI, MAI, All versions of MDS
Evaluating the overall diet quality in Mediterranean populations	MDS

Abbreviations

Not applicable

Declaration

Ethics approval and consent to participate

Ethics Committee of the university does not grant a code of ethics for review studies. However, the authors tried to follow all the ethical guidelines.

Availability of data and materials section

Not applicable

Consent for Publication

Not applicable

Financial disclosure

Authors have no competing interests to declare.

Funding/Support

This study did not receive funds from institutes or centers.

Authors' contributions

All authors have read and approved the manuscript.

Conceptualization RH Methodology: RH, SRH, FR Resources: SRH, FR, AR

Validation: JMR, RH

Writing (original draft preparation): SE

Investigation: SRH, FR **Visualization:** AH, FR, SRH

Writing (review and editing): JMR, MF, AR, SE

Project administration: RH

Supervision: RH

Acknowledgement

The authors thank all personnel at Noncommunicable Diseases Research Center (NCDRC) of Fasa University of Medical Sciences for their collaborations.

References

- Trichopoulou A, Vasilopoulou E. Mediterranean diet and longevity. British Journal of Nutrition. 2007;84(S2):S205-S9.
- Trichopoulou A. Traditional Mediterranean diet and longevity in the elderly: a review. Public Health Nutr. 2007;7(7):943-7.
- Pérez-López FR, Chedraui P, Haya J, Cuadros JL. Effects of the Mediterranean diet on longevity and age-related morbid conditions. Maturitas. 64(2):67-79.
- 4. Trichopoulou A, Critselis E. Mediterranean diet and longevity. European Journal of Cancer Prevention. 2004;13(5):453-6.
- Van Itallie TB, Yang M-U. Diet and Weight Loss. New England Journal of Medicine. 1977;297(21):1158-61.
- McManus K, Antinoro L, Sacks F. A randomized controlled trial of a moderate-fat, low-energy diet compared with a low fat, low-energy diet for weight loss in overweight adults. International Journal Of Obesity. 2001;25:1503.
- Ross R, Dagnone D, Jones PH, et al. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men: A randomized, controlled trial. Annals of Internal Medicine. 2000;133(2):92-103.
- 8. Kirk S. Diet and weight management. Nurs Stand. 2003;17(49):47-53; quiz 4-5.
- Zand H, Homayounfar R, Cheraghpour M, Jeddi-Tehrani M, Ghorbani A, Pourvali K, et al. Obesity-induced p53 activation in insulin-dependent and independent tissues is inhibited by beta-adrenergic agonist in diet-induced obese rats. Life sciences. 2016;147:103-9.
- Seely S, Freed DLJ, Silverstone GA, Rippere V. Diet-related diseases. The modern epidemic. Beckenham, Kent: Croom Helm Ltd.; 1985. 272pp. p.
- Diet, nutrition and the prevention of chronic diseases. Report of a WHO Study Group. Geneva: WHO; 1990.
- Ghaemi A, Hosseini N, Osati S, mehdi Naghizadeh M, Ehrampoush E, Honarvar B, et al. Waist circumference is a mediator of dietary pattern in Non-alcoholic fatty liver disease. 2018;8(1):4788.

- Sánchez-Villegas A, Henríquez P, Bes-Rastrollo M, Doreste J. Mediterranean diet and depression. Public Health Nutr. 2006;9(8A):1104-9.
- 14. Sanchez-Villegas A, Martínez-González MA. Diet, a new target to prevent depression? BMC Medicine. 2013;11(1):3.
- Quirk SE, Williams LJ, O'Neil A, Pasco JA, Jacka FN, Housden S, et al. The association between diet quality, dietary patterns and depression in adults: a systematic review. BMC Psychiatry. 2013;13(1):175.
- Felice NJ, Peter JK, Eva RL, Michael B, George CP, John WT, et al. Associations Between Diet Quality and Depressed Mood in Adolescents: Results from the Australian Healthy Neighbourhoods Study. Australian & New Zealand Journal of Psychiatry. 2010;44(5):435-42.
- 17. Skinner JD, Bounds W, Carruth BR, Morris M, Ziegler P. Predictors of children's body mass index: a longitudinal study of diet and growth in children aged 2–8 y. International Journal of Obesity. 2004;28(4):476-82.
- Rato L, Alves MG, Cavaco JE, Oliveira PF. High-energy diets: a threat for male fertility? Obesity Reviews. 2014;15(12):996-1007
- 19. Luck MR, Jeyaseelan I, Scholes RA. Ascorbic acid and fertility. Biology of Reproduction. 1995;52(2):262-6.
- 20. Myles IA. Fast food fever: reviewing the impacts of the Western diet on immunity. Nutrition Journal. 2014;13(1):61.
- Alkerwi A. Diet quality concept. Nutrition (Burbank, Los Angeles County, Calif). 2014;30(6):613-8.
- Preedy VR, Hunter LA, Patel VB. Diet Quality: An Evidence-Based Approach: Springer New York; 2013.
- 23. Diet quality 2017 [Available from: https://www.iaea.org/topics/diet-quality.
- 24. Wirt A, Collins CE. Diet quality--what is it and does it matter? Public Health Nutr. 2009;12(12):2473-92.
- 25. Patterson RE, Haines PS, Popkin BM. Diet quality index: Capturing a multidimensional behavior. Journal of the American Dietetic Association. 1994;94(1):57-64.
- Waijers PM, Feskens EJ, Ocké MC. A critical review of predefined diet quality scores. British Journal of Nutrition. 2007;97(2):219-31.
- Gil Á, Martinez de Victoria E, Olza J. Indicators for the evaluation of diet quality. Nutr Hosp [Internet]. 2015 2015/02//; 31 Suppl 3:[128-44 pp.]. Available from: http://europepmc.org/abstract/MED/25719781

https://doi.org/10.3305/nh.2015.31.sup3.8761.

- 28. R.D. L, D.C. N. Nutritional assessment. 2nd ed. 1996.
- 29. Haines PS, Siega-Riz AM, Popkin BM. The Diet Quality Index revised: a assessment instrument for populations. Journal of the American Dietetic Association. 1999;99(6):697-704.
- Newby P, Hu FB, Rimm EB, Smith-Warner SA, Feskanich D, Sampson L, et al. Reproducibility and validity of the Diet Quality Index Revised as assessed by use of a food-frequency questionnaire. The American Journal of Clinical Nutrition. 2003;78(5):941-9.
- 31. Fung TT, Hu FB, McCullough ML, Newby P, Willett WC, Holmes MD. Diet quality is associated with the risk of estrogen receptor–negative breast cancer in postmenopausal women. The Journal of nutrition. 2006;136(2):466-72.

Nutrition and Food Sciences Research

50

Vol 11, No 4, Oct-Dec 2024

- 32. Fung TT, McCullough ML, Newby P, Manson JE, Meigs JB, Rifai N, et al. Diet-quality scores and plasma concentrations of markers of inflammation and endothelial dysfunction—. The American journal of clinical nutrition. 2005;82(1):163-73.
- 33. Kim S, Haines PS, Siega-Riz AM, Popkin BM. The Diet Quality Index-International (DQI-I) provides an effective tool for cross-national comparison of diet quality as illustrated by China and the United States. The Journal of Nutrition. 2003;133(11):3476-84.
- 34. T Kennedy E, Ohls J, Carlson S, Fleming K. The Healthy Eating Index: Design and Applications. Journal of the American Dietetic Association. 1995;95(10):1103-8.
- 35. Osati S, Homayounfar R, Hajifaraji M. Metabolic effects of vitamin D supplementation in vitamin D deficient patients (a double-blind clinical trial). Diabetes & Metabolic Syndrome: Clinical Research & Reviews. 2016;10(2):S7-S10.
- 36. McCullough ML, Feskanich D, Stampfer MJ, Giovannucci EL, Rimm EB, Hu FB, et al. Diet quality and major chronic disease risk in men and women: moving toward improved dietary guidance. The American journal of clinical nutrition. 2002;76(6):1261-71.
- 37. Huijbregts P, Feskens E, Rasanen L, Fidanza F, Nissinen A, Menotti A, et al., editors. Dietary patterns and 20-year mortality in elderly men in Finland, Italy and the Netherlands. Perspectives on epidemiology in Europe Abstracts IEA Regional Meeting, The Hague (1995) 2205; 1995.
- 38. Huijbregts P, Feskens E, Räsänen L, Fidanza F, Alberti-Fidanza A, Nissinen A, et al. Dietary patterns and cognitive function in elderly men in Finland, Italy and The Netherlands. European Journal of Clinical Nutrition. 1998;52(11):826.
- 39. Osler M andreasen AH, Heitmann B, Høidrup S, Gerdes U, Jørgensen LM, et al. Food intake patterns and risk of coronary heart disease: a prospective cohort study examining the use of traditional scoring techniques. European journal of clinical nutrition. 2002;56(7):568.
- Osler M, Heitmann BL, Gerdes LU, Jørgensen LM, Schroll M. Dietary patterns and mortality in Danish men and women: a prospective observational study. British Journal of Nutrition. 2001;85(2):219-25.
- 41. Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, Gnardellis C, Lagiou P, Polychronopoulos E, et al. Diet and overall survival in elderly people. Bmj. 1995;311(7018):1457-60.

- 42. Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. New England Journal of Medicine. 2003;348(26):2599-608.
- 43. Osler M, Schroll M. Diet and mortality in a cohort of elderly people in a north European community. International journal of epidemiology. 1997;26(1):155-9.
- 44. Knoops K, Fidanza F, Alberti-Fidanza A, Kromhout D, Van Staveren W. Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. European journal of clinical nutrition. 2006;60(6):746.
- 45. Hatløy A, Torheim L, Oshaug A. Food variety—a good indicator of nutritional adequacy of the diet? A case study from an urban area in Mali, West Africa. European Journal of Clinical Nutrition. 1998;52(12):891.
- 46. Steyn N, Nel J, Nantel G, Kennedy G, Labadarios D. Food variety and dietary diversity scores in children: are they good indicators of dietary adequacy? Public Health Nutr. 2006;9(5):644-50.
- 47. Savy M, Martin-Prével Y, Sawadogo P, Kameli Y, Delpeuch F. Use of variety/diversity scores for diet quality assessment: relation with nutritional status of women in a rural area in Burkina Faso. European journal of clinical nutrition. 2005;59(5):703.
- 48. Kennedy GL, Pedro MR, Seghieri C, Nantel G, Brouwer I. Dietary diversity score is a useful indicator of micronutrient intake in non-breast-feeding Filipino children. The Journal of nutrition. 2007;137(2):472-7.
- Patrick MJ, Yoder MD. Program Evaluation: Food Stamps and Commodity Distribution in Rural Areas of Central Pennsylvania. 1971.
- 50. Seymour JD, Calle EE, Flagg EW, Coates RJ, Ford ES, Thun MJ. Diet Quality Index as a Predictor of Short-term Mortality in the American Cancer Society Cancer Prevention Study II Nutrition Cohort. American Journal of Epidemiology. 2003;157(11):980-8.
- 51. Fung TT, Hu FB, Mccullough ML, Newby P, Willett WC, Holmes MD. Diet quality is associated with the risk of estrogen receptor-negative breast cancer in postmenopausal women. The Journal of nutrition. 2006;136(2):466-72.